



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : Vol. ISSUE : 02 Print / Issue Publication Date: 09-Sep-2024
9, Issue 02



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2024.v09i02.017

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



A NOTE ON STRONG Σ^* – \mathcal{H} – OPEN SETS IN GTS VIA HEREDITARY CLASS

R. Shankar

Department Of Mathematics,
 Dwaraka Doss Goverdhan Doss Vaishnav College,
 Chennai, Tamil Nadu, India.

Abstract - The aim of this paper is to introduce and study the strong σ^* – \mathcal{H} – open set in a generalized topological space (X, λ) with a hereditary class \mathcal{H} . We give discuss and some of their properties strong σ^* – \mathcal{H} – open set and strong π^* – \mathcal{H} – open set in a generalized topological space (X, λ) with a hereditary class \mathcal{H} . Also discuss various characterization strong σ^* – \mathcal{H} – open set in a generalized topological space (X, λ) with a hereditary class \mathcal{H} are given and its properties of such sets are discussed. Also we establish state some characterize of strong π^* – \mathcal{H} – open set in a generalized topological space (X, λ) with a hereditary class \mathcal{H} .

Keywords - Hereditary class, π^* – \mathcal{H} – open set, λ – codense, τ – \mathcal{H} – set, σ^* – \mathcal{H} – open set, strong β – \mathcal{H} – open set and \mathcal{H} – open set.

2010 Mathematics subject classification: 54A05, 54A10.

I. INTRODUCTION

Let X be a nonempty set and $\wp(X)$ denotes the power set of X . Then the collection $\lambda \subseteq \wp(X)$ is called a generalized topology (GT) on X [1] if $\emptyset \in \lambda$ and arbitrary union of members of λ belongs to λ . A set X , with a GT on it is said to be generalized topological space (GTS) (X, λ) . Members of λ are called λ – open sets. A subset $E \subset X$ is said to be λ – closed sets if $X - E$ is λ – open set. For each $E \subset X$, we denote by $\text{int}_\lambda(E)$ [2] the union of all λ – open sets contained in E and by $\text{cl}_\lambda(E)$ [2] the intersection of all λ – closed sets containing E . A GTS (X, λ) is said to be a quasitopological space [4] (QTS) if $K, L \in \lambda$ implies $K \cap L \in \lambda$. A hereditary class (HC) \mathcal{H} is a nonempty family of subset of X such that $L \subset M, M \in \mathcal{H}$ implies $L \in \mathcal{H}$ [2]. Hereditary class \mathcal{H} is said to be λ – codense [2] if $\lambda \cap \mathcal{H} = \{\emptyset\}$ [2]. For each $E \subset X$, a subset $E^*(\mathcal{H})$ is defined by $E^* = \{x \in X \mid N \cap E \notin \mathcal{H} \text{ for every } N \in \lambda \text{ such that } x \in N\}$ [3]. If $\text{cl}_\lambda^*(E) = E \cup E^*$ for each $E \subset X$, with respect to λ and a HC \mathcal{H} of subsets of X , then $\lambda^* = \{E \subset X \mid \text{cl}_\lambda^*(X - E) = X - E\}$ [3] is a GT. Members of λ^* are called λ^* – open sets and its complement is called a λ^* – closed set. We denote the interior of E in (X, λ^*) is $\text{int}_{\lambda^*}(E)$. A subset $E \subset X$ is said to be λ^* – dense (resp. λ – dense) if $\text{cl}_{\lambda^*}(E) = X$ (resp. $\text{cl}_\lambda(E) = X$).

Literature Review

In 1997, Prof. Csaszar [1] nicely pointed out that the family of all open sets and all weak forms of open sets in a topological space (X, τ) can be obtained from monotonic functions defined on $\wp(X)$, the family of all subsets of X . His further study of generalized open sets using monotonic functions on $\wp(X)$. The collection of all monotonic functions from $\wp(X)$ to $\wp(X)$ where X is any non-empty set and denote it by $\Gamma(X)$. For $\gamma \in \Gamma, \lambda = \{E \subset X \mid E \subset \gamma(E)\}$. In 2002, Csaszar, defined generalized topology [2]. He proved that $\emptyset \in \lambda$ and λ is closed under arbitrary union. He named such families as Generalized Topology and the pair (X, λ) is called generalized topological space. Elements of λ are called λ – open sets and the complement of an λ – open set is called a λ – closed set. The union of all λ – open sets contained in a subset E of X is denoted by $\text{int}_\lambda(E)$ and is called the λ – interior of E . The intersection of all λ – closed sets containing E is called the λ – closure of E and is denoted by $\text{cl}_\lambda(E)$. In 2005, Csaszar [2] defined the generalized open sets and discuss their relationship for a generalized topology λ on a nonempty set X , the generalized open sets λ – α – open, λ – σ – open, λ – π – open and λ – β – open sets. In 2006, Csaszar defined the normal generalized topological space. In 2007, Csaszar [3] defined a nonempty class of subsets of a nonempty set X , called as hereditary class \mathcal{H} and studied modification of generalized topology with hereditary classes. He defined the concepts codense and strongly completely codense hereditary classes. Also he defined about quazi topology, which is closed under finite intersection. In 2009, Csaszar defined and studied the product of generalized topologies. Also, generalized topology with hereditary classes are studied by Y.K. Kim and W.K. Min [21], V. Renukadevi and Sheena Scaria [5], V. Renukadevi and K. Karuppaiy [6].

II. MATERIALS AND METHODS

Lemma 2.1 [1] Let (X, λ) be a GTS and $E, F \subset X$. Then the following hold.

- (i) $\text{int}_\lambda(\emptyset) = \emptyset$.
- (ii) $\text{int}_\lambda(X) = X$.
- (iii) $\text{int}_\lambda(\text{int}_\lambda(E)) = \text{int}_\lambda(E)$.
- (iv) $E \in \lambda \Leftrightarrow \text{int}_\lambda(E) = E$.
- (v) $\text{int}_\lambda(E \cap F) = \text{int}_\lambda(E) \cap \text{int}_\lambda(F)$.



(vi) $\text{int}_\lambda(E) \cup \text{int}_\lambda(F) \subset \text{int}_\lambda(E \cup F)$.

Lemma 2.2 [1] Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E, F \subset X$. Then the following hold.

- (i) $\text{cl}_\lambda(\emptyset) = \emptyset$
- (ii) $\text{cl}_\lambda(X) = X$.
- (iii) E is λ -closed $\Leftrightarrow \text{cl}_\lambda(E) = E$.
- (iv) $\text{cl}_\lambda(E \cap F) \subset \text{cl}_\lambda(E) \cap \text{cl}_\lambda(F)$.
- (v) $\text{cl}_\lambda(E \cup F) = \text{cl}_\lambda(E) \cup \text{cl}_\lambda(F)$
- (vi) $\text{cl}_\lambda(\text{cl}_\lambda(E)) = \text{cl}_\lambda(E)$.

Lemma 2.3. [3] Let (X, λ) be a GTS with a HC \mathcal{H} and $S, T \subset X$. Then the following hold.

- (i) $S \subset T$ implies $S^* \subset T^*$.
- (ii) $(S^*)^* = S^*$ for every $S \subset X$.
- (iii) $S \subset T \subset X$ implies $\text{cl}_\lambda^*(S) \subset \text{cl}_\lambda^*(T)$.
- (iv) $(E \cup E^*)^* \subset E^*$ for every $E \subset X$.
- (v) $(S \cup T)^* = S^* \cup T^*$.
- (vi) $\lambda \subset \lambda^*$.
- (vii) $\text{Fis}\lambda^*$ -closed if and only if $F^* \subset F$.
- (viii) $\beta = \{N - K : N \in \lambda, K \in \mathcal{H}\}$ is a base for λ^* .

Lemma 2.4. [6] Let (X, λ) be a GTS with a HC \mathcal{H} and $E \subset X$. Then the following hold.

- (i) If $N \in \lambda$, then $N \cap \text{cl}_\lambda^*(E) \subset \text{cl}_\lambda^*(N \cap E)$.
- (ii) If $N \in \lambda$, then $N \cap E^* \subset (N \cap E)^*$.

Lemma 2.5.[5] Let (X, λ) be a GTS with a λ -codense hereditary class \mathcal{H} and $E \subset X$. If $E \subset E^*$, then $E^* = \text{cl}_\lambda(E) = \text{cl}_\lambda^*(E) = \text{cl}_\lambda(E^*)$.

Lemma 2.6.[3] Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. If E is a λ -semi closed set, then $\text{int}_\lambda(E) = \text{int}_\lambda(\text{cl}_\lambda(E))$.

Lemma 2.7. [21] Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. Then the following hold.

- (a) $\text{int}_\lambda(E) \subset \text{int}_\lambda^*(E) \subset E$.
- (b) $\text{int}_\lambda^*(E) = X - \text{cl}_\lambda^*(X - E)$.
- (c) $\text{cl}_\lambda^*(E) = X - \text{int}_\lambda^*(X - E)$.
- (d) $E \subset \text{cl}_\lambda^*(E) \subset \text{cl}_\lambda(E)$.

Lemma 2.8.[21] Let (X, λ) be a GTS with a λ -codense hereditary class \mathcal{H} and $E \subset X$. Then the following hold.

- (i) $\text{int}_\lambda(E) = \text{int}_\lambda^*(E)$, for every λ^* -closed set E .
- (ii) $\text{cl}_\lambda(E) = \text{cl}_\lambda^*(E)$, for every λ^* -open set E .


Definition 2.9. A subset $E \subset X$ of a GTS (X, λ) with a hereditary class \mathcal{H} is said to be

- 1. λ^* -Dense [3] in itself if $E \subset E^*$.
- 2. λ^* -Perfect [3] if $E = E^*$.
- 3. λ^* -Closed [3] if $E^* \subset E$.
- 4. \mathcal{H} -open [3] if $E \subset \text{int}_\lambda(E^*)$

- 5. $\sigma - \mathcal{H}$ -open [3] if $E \subset \text{cl}_\lambda^*(\text{int}_\lambda(E))$.
- 6. Weakly semi- \mathcal{H} -open [11] if $E \subset \text{cl}_\lambda^*(\text{int}_\lambda(\text{cl}_\lambda(E)))$.
- 7. Almost strong- \mathcal{H} -open [11] if $E \subset \text{cl}_\lambda^*(\text{int}_\lambda(E^*))$.
- 8. $\sigma^* - \mathcal{H}$ -open [6] if $E \subset \text{cl}_\lambda(\text{int}_\lambda^*(E))$.
- 9. $\pi^* - \mathcal{H}$ -open [6] if $E \subset \text{int}_\lambda^*(\text{cl}_\lambda(E))$.
- 10. $\beta^* - \mathcal{H}$ -open [6] if $E \subset \text{cl}_\lambda(\text{int}_\lambda^*(\text{cl}_\lambda(E)))$.
- 11. $\pi - \mathcal{H}$ -open [3] if $E \subset \text{int}_\lambda(\text{cl}_\lambda^*(E))$.
- 12. $\alpha - \mathcal{H}$ -open [3] if $E \subset \text{int}_\lambda(\text{cl}_\lambda^*(\text{int}_\lambda(E)))$.
- 13. $\mathcal{H}\mathcal{R}$ -closed set [12] $E = \text{cl}_\lambda^*(\text{int}_\lambda(E))$.
- 14. $\alpha^* - \mathcal{H}$ -open [16] if $\text{int}_\lambda(\text{cl}_\lambda^*(\text{int}_\lambda(E))) = \text{int}_\lambda(E)$.
- 15. $t - \mathcal{H}$ -set [7] if $\text{int}_\lambda(E) = \text{int}_\lambda(\text{cl}_\lambda^*(E))$.
- 16. Strong $\beta - \mathcal{H}$ -open [13] if $E \subset \text{cl}_\lambda^*(\text{int}_\lambda(\text{cl}_\lambda^*(E)))$.
- 17. Strong $\pi^* - \mathcal{H}$ -open [22] set if $\text{int}_\lambda^*(\text{cl}_\lambda^*(E))$.

The complement of a strong $\beta - \mathcal{H}$ -open (resp. $\sigma - \mathcal{H}$ -open, $\sigma^* - \mathcal{H}$ -open, $\pi^* - \mathcal{H}$ -open, $\beta^* - \mathcal{H}$ -open, $\alpha^* - \mathcal{H}$ -open, $\pi - \mathcal{H}$ -open, weakly semi- \mathcal{H} -open, Strong $\pi^* - \mathcal{H}$ -open, $\alpha - \mathcal{H}$ -open) set is said to a strong $\beta - \mathcal{H}$ -closed (resp. $\sigma - \mathcal{H}$ -Closed, $\sigma^* - \mathcal{H}$ -closed, $\pi^* - \mathcal{H}$ -closed, $\beta^* - \mathcal{H}$ -closed, $\alpha^* - \mathcal{H}$ -closed, $\pi - \mathcal{H}$ -closed, weakly semi- \mathcal{H} -closed, Strong $\pi^* - \mathcal{H}$ -closed, $\pi - \mathcal{H}$ -closed, $\alpha - \mathcal{H}$ -closed) set.

III. RESULTS AND DISCUSSION



3.1 Strong $\sigma^* - \mathcal{H}$ -open Sets

Definition 3.1. A subset $E \subset X$ of a GTS (X, λ) with a hereditary class \mathcal{H} is said to be a strong $\sigma^* - \mathcal{H}$ -open set if $E \subset \text{cl}_\lambda^*(\text{int}_\lambda^*(E))$. The family of all strong $\sigma^* - \mathcal{H}$ -open set is denoted by $S\sigma^*\mathcal{H}(X, \lambda)$. If its complement is strong $\sigma^* - \mathcal{H}$ -closed set.



Theorem 3.2. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. If every $\sigma - \mathcal{H}$ -open set is a strong $\sigma^* - \mathcal{H}$ -open set.

Proof. Let E be a $\sigma - \mathcal{H}$ -open set. Then $E \subset \text{cl}_\lambda^*(\text{int}_\lambda(E))$. Since λ^* finer than λ and so $E \subset \text{cl}_\lambda^*(\text{int}_\lambda(E)) \subset \text{cl}_\lambda^*(\text{int}_\lambda^*(E))$. Therefore, E is a strong $\sigma^* - \mathcal{H}$ -open set. The following Example 3.3 shows that the converse of Theorem 3.2 is not true.

Example 3.3. Let

$X = \{\chi_1, \chi_2, \chi_3, \chi_4\}, \lambda = \{\emptyset, X, \{\chi_1\}, \{\chi_1, \chi_2\}, \{\chi_3, \chi_4\}, \{\chi_1, \chi_3, \chi_4\}\}$ and $\mathcal{H} = \{\emptyset, \{\chi_1\}, \{\chi_4\}, \{\chi_1, \chi_4\}\}$. If $E = \{\chi_3\}$ then $E^* = \{\chi_2, \chi_3, \chi_4\}$. Now $\text{cl}_\lambda^*(\text{int}_\lambda^*(E)) = \text{cl}_\lambda^*(\text{int}_\lambda^*(\{\chi_3\})) = \text{cl}_\lambda^*(X - \text{cl}_\lambda^*(X - \{\chi_3\})) = \text{cl}_\lambda^*(X - \text{cl}_\lambda^*(\{\chi_1, \chi_2, \chi_4\})) = \text{cl}_\lambda^*(X - (\{\chi_1, \chi_2, \chi_4\} \cup \{\chi_2\})) = \text{cl}_\lambda^*(X - \{\chi_1, \chi_2, \chi_4\}) = \text{cl}_\lambda^*(\{\chi_3\}) = \{\chi_3\} \cup \{\chi_2, \chi_3, \chi_4\} = \{\chi_2, \chi_3, \chi_4\} \supset E$. Therefore E is a strong $\sigma^* - \mathcal{H}$ -open set. But $\text{cl}_\lambda^*(\text{int}_\lambda(E)) = \text{cl}_\lambda^*(\emptyset) = \emptyset \not\supset E$. Hence E is not a $\sigma - \mathcal{H}$ -open set.

Theorem 3.4. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. If every strong $\sigma^* - \mathcal{H}$ -open set is a $\sigma - \mathcal{H}$ -open set.

Proof. Let E be a strong $\sigma^* - \mathcal{H}$ -open set. Then $E \subset \text{cl}_\lambda^*(\text{int}_\lambda^*(E))$. Since λ^* finer than λ and so $E \subset \text{cl}_\lambda^*(\text{int}_\lambda^*(E)) \subset \text{cl}_\lambda(\text{int}_\lambda^*(E))$. Therefore, E is a $\sigma - \mathcal{H}$ -open set. The following Example 3.5 shows that the converse of Theorem 3.4 is not true.

Example 3.5. Let

$X = \{\chi_1, \chi_2, \chi_3, \chi_4\}, \lambda = \{\emptyset, X, \{\chi_1\}, \{\chi_2\}, \{\chi_1, \chi_2\}\}$ and $\mathcal{H} = \{\emptyset, \{\chi_2\}, \{\chi_3\}, \{\chi_2, \chi_3\}\}$. If $E = \{\chi_2, \chi_3\}$ then $E^* = \emptyset$. Now, $\text{cl}_\lambda(\text{int}_\lambda^*(E)) = \text{cl}_\lambda(\text{int}_\lambda^*(\{\chi_2, \chi_3\})) = \text{cl}_\lambda(X - \text{cl}_\lambda^*(X - \{\chi_2, \chi_3\})) = \text{cl}_\lambda(X - \text{cl}_\lambda^*(\{\chi_1, \chi_4\})) = \text{cl}_\lambda(X - (\{\chi_1, \chi_4\} \cup \{\chi_1, \chi_3, \chi_4\})) = \text{cl}_\lambda(X - \{\chi_1, \chi_3, \chi_4\}) = \text{cl}_\lambda(\{\chi_2\}) = \{\chi_2, \chi_3, \chi_4\} \supset E$. Therefore E is a strong $\sigma^* - \mathcal{H}$ -open set. But $\text{cl}_\lambda^*(\text{int}_\lambda(E)) = \text{cl}_\lambda^*(\text{int}_\lambda(\{\chi_2, \chi_3\})) = \text{cl}_\lambda^*(X - \text{cl}_\lambda^*(X - \{\chi_2, \chi_3\})) = \text{cl}_\lambda^*(X - \text{cl}_\lambda^*(\{\chi_1, \chi_4\})) = \text{cl}_\lambda^*(X - \{\chi_1, \chi_3, \chi_4\}) = \text{cl}_\lambda^*(\{\chi_2\}) = \{\chi_2\} \cup \emptyset = \{\chi_2\} \not\supset E$. Hence E is not a strong $\sigma - \mathcal{H}$ -open set.

Theorem 3.6. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. Then E is a strong $\sigma^* - \mathcal{H}$ -open set $\Leftrightarrow \text{cl}_\lambda^*(E) = \text{cl}_\lambda^*(\text{int}_\lambda^*(E))$.

Proof. Let E be a strong $\sigma^* - \mathcal{H}$ -open set. Then $E \subset \text{cl}_\lambda^*(\text{int}_\lambda^*(E))$. By Lemma, $\text{cl}_\lambda^*(E) \subset \text{cl}_\lambda^*(\text{cl}_\lambda^*(\text{int}_\lambda^*(E))) = \text{cl}_\lambda^*(\text{int}_\lambda^*(E)) \subset \text{cl}_\lambda^*(E)$. Hence $\text{cl}_\lambda^*(E) = \text{cl}_\lambda^*(\text{int}_\lambda^*(E))$.

Conversely, let $\text{cl}_\lambda^*(E) = \text{cl}_\lambda^*(\text{int}_\lambda^*(E))$, since $E \subset \text{cl}_\lambda^*(E) \Rightarrow E \subset \text{cl}_\lambda^*(\text{int}_\lambda^*(E))$.

Theorem 3.7. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. If E is a $\mathcal{H}R$ -closed set then, E is a strong $\sigma^* - \mathcal{H}$ -open set.

Proof. Let E be a $\mathcal{H}R$ -closed set. Then $E = \text{cl}_\lambda^*(\text{int}_\lambda(E)) \subset \text{cl}_\lambda^*(\text{int}_\lambda^*(E)) \Rightarrow E$ is a strong $\sigma^* - \mathcal{H}$ -open set.

The following Example 3.8 shows that the converse of Theorem 3.7 is not true.

Example 3.8. Let

$X = \{\chi_1, \chi_2, \chi_3, \chi_4\}, \lambda = \{\emptyset, X, \{\chi_4\}, \{\chi_1, \chi_3\}, \{\chi_1, \chi_3, \chi_4\}\}$ and $\mathcal{H} = \{\emptyset, \{\chi_3\}, \{\chi_4\}, \{\chi_3, \chi_4\}\}$. If $E = \{\chi_1\}$. Now $\text{cl}_\lambda^*(\text{int}_\lambda^*(E)) = \text{cl}_\lambda^*(X - \text{cl}_\lambda^*(X - E)) = \text{cl}_\lambda^*(X - \text{cl}_\lambda^*(\{\chi_2, \chi_3, \chi_4\})) = \text{cl}_\lambda^*(X - \{\chi_2, \chi_3, \chi_4\}) = \text{cl}_\lambda^*(\{\chi_1\}) = \{\chi_1, \chi_2, \chi_3\} \supset E$. Therefore E is a strong $\sigma^* - \mathcal{H}$ -open set. But $\text{cl}_\lambda^*(\text{int}_\lambda(E)) = \text{cl}_\lambda^*(\text{int}_\lambda(\{\chi_1\})) = \text{cl}_\lambda^*(\emptyset) = \emptyset \neq E$. Hence E is not a $\mathcal{H}R$ -closed set.

Theorem 3.9. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. If E is a strong $\pi^* - \mathcal{H}$ -open set, then $\text{cl}_\lambda^*(E)$ is a strong $\sigma^* - \mathcal{H}$ -open set.

Proof. Let E be a strong $\pi^* - \mathcal{H}$ -open set. Then, $E \subset \text{int}_\lambda^*(\text{cl}_\lambda^*(E))$, by Lemma $\text{cl}_\lambda^*(E) \subset \text{cl}_\lambda^*(\text{int}_\lambda^*(\text{cl}_\lambda^*(E))) \Rightarrow \text{cl}_\lambda^*(E)$ is a strong $\sigma^* - \mathcal{H}$ -open set.

Theorem 3.10. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. If E is a strong $\sigma^* - \mathcal{H}$ -open set, then $\text{int}_\lambda^*(E)$ is a strong $\pi^* - \mathcal{H}$ -open set.

Proof. Let E be a strong $\sigma^* - \mathcal{H}$ -open set. Then $E \subset \text{cl}_\lambda^*(\text{int}_\lambda^*(E))$, by Lemma $\text{int}_\lambda^*(E) \subset \text{int}_\lambda^*(\text{cl}_\lambda^*(\text{int}_\lambda^*(E))) \Rightarrow \text{int}_\lambda^*(E)$ is a $\pi^* - \mathcal{H}$ -open set.

Theorem 3.11. Let (X, λ) be a GTS with a λ -codense hereditary class \mathcal{H} and $E \subset X$. Then E is a $\alpha - \mathcal{H}$ -open set $\Leftrightarrow E$ is a strong $\sigma^* - \mathcal{H}$ -open set and strong $\pi^* - \mathcal{H}$ -open set.

Proof. Let E be a $\alpha - \mathcal{H}$ -open set. Since every $\alpha - \mathcal{H}$ -open set is both a $\sigma - \mathcal{H}$ -open set and $\pi - \mathcal{H}$ -open set. Also, every $\sigma - \mathcal{H}$ -open is a strong $\sigma^* - \mathcal{H}$ -open set and every $\pi - \mathcal{H}$ -open set is a strong $\pi^* - \mathcal{H}$ -open set. Conversely, Suppose E is a strong $\sigma^* - \mathcal{H}$ -open set and strong $\pi^* - \mathcal{H}$ -open set. Then we have, $E \subset \text{int}_\lambda^*(\text{cl}_\lambda^*(E)) \subset \text{int}_\lambda^*(\text{cl}_\lambda^*(\text{cl}_\lambda^*(\text{int}_\lambda^*(E)))) = \text{int}_\lambda^*(\text{cl}_\lambda^*(\text{int}_\lambda^*(E))) = \text{int}_\lambda(\text{cl}_\lambda^*(\text{int}_\lambda(E)))$. Hence E is a $\alpha - \mathcal{H}$ -open set.



Theorem 3.12. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$ and E is $\lambda - \pi$ -open set. If E is a semi closed set, then E is a strong $\sigma^* - \mathcal{H}$ -open set.

Proof. Suppose E is a semi closed set. Then, $int_\lambda(cl_\lambda(E)) = int_\lambda(E)$. Since E is $\lambda - \pi$ -open set. Then, we have $E \subset int_\lambda(cl_\lambda(E)) = int_\lambda(E) \subset int_\lambda(E) \cup int_\lambda^*(E) = cl_\lambda^*(int_\lambda(E)) \subset cl_\lambda^*(int_\lambda^*(E))$. Therefore E is a strong $\sigma^* - \mathcal{H}$ -open set.

Theorem 3.13. Let (X, λ) be a GTS with λ -codense hereditary class \mathcal{H} and $E \subset X$ be a λ^* -dense in-itself. If E is a strong $\sigma^* - \mathcal{H}$ -open set, then E is a strong $\beta - \mathcal{H}$ -open set.

Proof. Let E be a strong $\sigma^* - \mathcal{H}$ -open set. Then $E \subset cl_\lambda^*(int_\lambda^*(E)) \subset cl_\lambda^*(int_\lambda^*(E^*)) = cl_\lambda^*(int_\lambda^*(cl_\lambda^*(E)))$. There E is a strong $\beta - \mathcal{H}$ -open set.

Theorem 3.14. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$ and E is λ^* -perfect set. If E is almost strong- \mathcal{H} -open set, then E is a strong $\sigma^* - \mathcal{H}$ -open set.

Proof. Let E is a almost strong \mathcal{H} -open set. Then $E \subset cl_\lambda^*(int_\lambda(E^*))$. Since E is λ^* -perfect set, we have $E = E^* \Rightarrow E \subset cl_\lambda^*(int_\lambda(E^*)) = cl_\lambda^*(int_\lambda(E)) \subset cl_\lambda^*(int_\lambda^*(E))$. Therefore E is a strong $\sigma^* - \mathcal{H}$ -open set.

Theorem 3.15. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$ and E is λ^* -perfect set. If E is \mathcal{H} -open set, then E is a strong $\sigma^* - \mathcal{H}$ -open set.

Proof. Let E is a \mathcal{H} -open set. Then $E \subset int_\lambda(E^*)$. Since E is a λ^* -perfect set and λ^* is finer the λ , $E \subset int_\lambda(E) \subset cl_\lambda^*(int_\lambda(E)) \subset cl_\lambda^*(int_\lambda^*(E))$. Hence E is a strong $\sigma^* - \mathcal{H}$ -open set.

Theorem 3.16. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E, F \subset X$. If E is a strong $\sigma^* - \mathcal{H}$ -open set and F is a $\lambda - \sigma$ -open set, then $E \cup F$ is a $\sigma^* - \mathcal{H}$ -open set.

Proof. Let E be a strong $\pi^* - \mathcal{H}$ -open set and F be a $\lambda - \sigma$ -open set. Then $E \subset cl_\lambda^*(int_\lambda^*(E))$ and $F \subset cl_\lambda(int_\lambda(F))$. Since λ^* finer than λ , now $E \cup F \subset cl_\lambda^*(int_\lambda^*(E)) \cup cl_\lambda(int_\lambda(F)) \subset cl_\lambda(int_\lambda^*(E)) \cup cl_\lambda(int_\lambda^*(F)) \subset cl_\lambda((int_\lambda^*(E)) \cup (int_\lambda^*(F))) \subset cl_\lambda(int_\lambda^*(E \cup F))$. Hence $E \cup F$ is a $\sigma^* - \mathcal{H}$ -open set.

Theorem 3.17. Let (X, λ) be a QTS with a hereditary class \mathcal{H} and $E, F \subset X$. If E is a strong $\sigma^* - \mathcal{H}$ -open set and F is a $\lambda - \sigma$ -open set, then $E \cap F$ is an strong $\sigma^* - \mathcal{H}$ -open set.

Proof. Let E be a strong $\sigma^* - \mathcal{H}$ -open set and F be a $\lambda - \sigma$ -open set. We have $E \subset cl_\lambda^*(int_\lambda^*(E))$ and $F = int_\lambda(F)$. Now, $E \cap F \subset cl_\lambda^*(int_\lambda^*(E)) \cap int_\lambda(F) = [int_\lambda^*(E) \cup$

$(int_\lambda^*(E))^*] \cap int_\lambda(F) = [int_\lambda^*(E) \cap int_\lambda(F)] \cup [(int_\lambda^*(E))^* \cap int_\lambda(F)] \subset [int_\lambda^*(E) \cap int_\lambda^*(F)] \cup [(int_\lambda^*(E))^* \cap int_\lambda^*(F)] \subset [int_\lambda^*(E \cap F)] \cup [int_\lambda^*(E) \cap int_\lambda^*(F)]^* \subset [int_\lambda^*(E \cap F)] \cup [int_\lambda^*(E \cap F)]^* = cl_\lambda^*(int_\lambda^*(E \cap F))$. Therefore, $E \cap F$ is a strong $\sigma^* - \mathcal{H}$ -open set.

Theorem 3.18. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. Then E is Strong $\sigma^* - \mathcal{H}$ -closed $\Leftrightarrow int_\lambda^*(cl_\lambda^*(E)) \subset E$.

Proof. Let E be a Strong $\sigma^* - \mathcal{H}$ -closed $\Leftrightarrow X - E$ is a strong $\sigma^* - \mathcal{H}$ -open set $\Leftrightarrow (X - E) \subset cl_\lambda^*(int_\lambda^*(X - E)) = cl_\lambda^*(X - cl_\lambda^*(E)) = X - int_\lambda^*(cl_\lambda^*(E)) \Leftrightarrow int_\lambda^*(cl_\lambda^*(E)) \subset E$.

Theorem 3.19. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E, F \subset X$. If E is a strong $\sigma^* - \mathcal{H}$ -closed set and F is a weakly semi- \mathcal{H} -closed set, then $E \cap F$ is a $\beta^* - \mathcal{H}$ -closed set.

Proof. Let E be a strong $\pi^* - \mathcal{H}$ -closed set and F be a weakly semi- \mathcal{H} -closed set. Then $E \supset int_\lambda^*(cl_\lambda^*(E))$ and $F \supset int_\lambda^*(cl_\lambda(int_\lambda(F)))$.

Now, $E \cap F \supset int_\lambda^*(cl_\lambda^*(E)) \cap int_\lambda^*(cl_\lambda(int_\lambda(F)))$ by Lemma,

$E \cap F \supset int_\lambda^*(cl_\lambda^*(int_\lambda(E))) \cap int_\lambda^*(cl_\lambda(int_\lambda(F))) = int_\lambda^*(cl_\lambda^*(int_\lambda(E)) \cap cl_\lambda^*(int_\lambda(F))) \supset int_\lambda^*(cl_\lambda^*(int_\lambda(E) \cap int_\lambda(F))) = int_\lambda^*(cl_\lambda^*(int_\lambda(E \cap F)))$. Hence $E \cap F$ is a $\beta^* - \mathcal{H}$ -closed set.

Theorem 3.20. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E, F \subset X$. If E is a strong $\sigma^* - \mathcal{H}$ -closed set and F is a $t - \mathcal{H}$ -set, then $E \cap F$ is a $\sigma^* - \mathcal{H}$ -closed set.

Proof. Let E be a strong $\pi^* - \mathcal{H}$ -closed set and F is a $t - \mathcal{H}$ -set. Then $E \supset int_\lambda^*(cl_\lambda^*(E))$ and $int_\lambda(F) = int_\lambda(cl_\lambda^*(F))$. Now, $E \cap F \supset int_\lambda^*(cl_\lambda^*(E)) \cap F \supset int_\lambda^*(cl_\lambda^*(E)) \cap int_\lambda(F) \supset int_\lambda^*(cl_\lambda^*(E)) \cap int_\lambda^*(cl_\lambda^*(F)) \supset int_\lambda^*(cl_\lambda^*(E)) \cap int_\lambda^*(cl_\lambda^*(F)) = int_\lambda^*(cl_\lambda^*(E) \cap cl_\lambda^*(F)) \supset int_\lambda^*(cl_\lambda^*(E \cap F))$. Therefore E is a $\sigma^* - \mathcal{H}$ -closed set.

Theorem 3.21. Let (X, λ) be a GTS with a hereditary class \mathcal{H} and $E \subset X$. Then E is a strong $\sigma^* - \mathcal{H}$ -open set \Leftrightarrow there exists a strong $\sigma^* - \mathcal{H}$ -open set F such that $int_\lambda^*(F) \subset E \subset F$.

Proof. Let E be a strong $\pi^* - \mathcal{H}$ -closed set. Then $E \supset int_\lambda^*(cl_\lambda^*(E))$. Let us assume $F = cl_\lambda^*(E)$ be a λ^* -closed set. There exists a strong $\sigma^* - \mathcal{H}$ -open set F such that by Lemma, $int_\lambda^*(F) = int_\lambda^*(cl_\lambda^*(E)) \subset E \subset cl_\lambda^*(E) = F$. Conversely, If F is a strong $\sigma^* - \mathcal{H}$ -closed



set such that $int_{\lambda}^*(F) \subset E \subset F$. Then by Lemma, $int_{\lambda}^*(E) = int_{\lambda}^*(F)$. Since F is a strong $\sigma^* - \mathcal{H}$ -closed set, $\Rightarrow int_{\lambda}^*(cl_{\lambda}^*(F)) \subset F$ and hence $E \supset int_{\lambda}^*(F) \supset int_{\lambda}^*(int_{\lambda}^*(cl_{\lambda}^*(F))) = int_{\lambda}^*(cl_{\lambda}^*(F)) = int_{\lambda}^*(cl_{\lambda}^*(E))$ it follows that, $E \supset int_{\lambda}^*(cl_{\lambda}^*(E))$. Therefore E is a strong $\sigma^* - \mathcal{H}$ -closed set.

IV. CONCLUSION

As stated in the abstract, we have defined the strong $\sigma^* - \mathcal{H}$ -open set in a Generalized Topological space (X, λ) with a hereditary class \mathcal{H} . Also, we discuss its properties and give various characterization of this sets. A definition of a strong $\sigma^* - \mathcal{H}$ -open set was defined, it has been shown that the concept of a strong $\sigma^* - \mathcal{H}$ -open set is weaker than the concept of a $\sigma - \mathcal{H}$ -open set and stronger than the concept of $\sigma^* - \mathcal{H}$ -open set. A discussion investigate the relationship between strong $\sigma^* - \mathcal{H}$ -open set and some known concepts Generalized Topological space (X, λ) with a hereditary class \mathcal{H} . Moreover, many counter examples were established.

V. ACKNOWLEDGEMENTS

The authors would like to express their sincere thanks to editors and the anonymous referees for their valuable comments and suggestions which improved the research article.

REFERENCES

- [1]. CsaszarA.(1997), Generalized Open Sets, Acta Math. Hungar, 75(1-2), (pp 65-87).
- [2]. CsaszarA. (2002), Generalized topology, generalized continuity, Acta Math. Hungar, 96, (pp 351-357).
- [3]. CsaszarA.(2007), Modifications of generalized topologies via hereditary classes, Acta Math. Hungar, 115,(pp 29 – 36).
- [4]. CsaszarA. (2007), Remarks on quasi-topologies, Acta Math. Hungar, 119 (1-2), (pp 197 – 200).
- [5]. RenukadeviV., and Sheena Scaria. (2011), On hereditary classes in generalized topological spaces, J. Adv. Res. Pure. Math., 3(2), (pp 21-30).
- [6]. RenukadeviV., and KaruppaiyK.(2010), On modifications of generalized topologies via hereditary classes, J. Adv. Res. Pure. Math., no.2. (pp 14-20).
- [7]. KaruppaiyK. (2013), Some subsets of GTS with hereditary classes, Journal of Advanced Studies in Topology. 5 (1), (pp 25-33).
- [8]. KaruppaiyK.(2014), A Note on $\delta - \mathcal{H}$ -sets in GTS with hereditary classes, International Journal of Mathematical Archive. 5(1), (pp 226-229).
- [9]. K.Karuppaiy, A Note on RH-open sets in GTS with hereditary classes, International Journal of Mathematical Archive. 5(1), (2014), 312-316.
- [10]. ShankarR., PonrajA.P.,and SivarajD.(2017), Weakly semi- \mathcal{H} -open sets and almost strong \mathcal{H} - κ -open sets, International Journal of Pure and Applied Mathematics. Vol-117 (17), (pp 7-13).
- [11]. PonrajA.P., ShankarR., and SivarajD. (2018), \mathcal{H}_R -closed set in generalized topological space, International Journal of Pure and Applied Mathematics. Vol-119(15), (pp 2153-2158).
- [12]. ShankarR., PonrajA.P., and SivarajD (2018), $A_{\mathcal{H}R}$ -sets in generalized topological space, International Journal of Pure and Applied Mathematics. Vol-119(15), (pp 2165-2170).
- [13]. ShankarR., and SivarajD. (2021), Strong $\beta - \mathcal{H}$ -Open sets in Generalized Topological Space, International Journal of Mathematics Trends and Technology, 67(7),(pp 183-186).
- [14]. RenukadeviV., and VimaladeviP.(2014), Note on generalized topological space with hereditary classes, Bolo. Soc. Paran. Mat. 32 (1), (pp 89-97).
- [15]. PankajamV., and SivarajD.(2013), Some separation axioms in GTS, Bolo. Soc. Paran. Mat. 31 (1), (pp 29-42).
- [16]. RenukadeviV., and SivarajD.(2014), Weak separation axioms in GTS, KYUNGPOOK Math, J. (54), 387-399.
- [17]. KaruppaiyK.(2014), On decomposition of continuity and complete continuity with hereditary class. J.Adv. Stud. Topology. Vol-(5), (pp.18-26).
- [18]. PonrajA.P., and SivarajD.(2017), $semi^* - \mathcal{H}$ -open sets, International Journal pure and applied mathematics. Vol-117(17), (pp 1-6).
- [19]. RenukadeviV., and VimaladeviP.(2015), Compatible Hereditary classes in GTS, Acta Mathematica Hungaric, Vol-147(2),(pp 259-271).
- [20]. UmitKarabiyik., and AynurKeskinKaymakci.(2014), On Semi- \mathcal{H}_S -open sets in Hereditary Generalized Topological Spaces, International Journal of Mathematical Analysis. Vol-8(53), (pp 2629-2637).
- [21]. Irina Zvina. (2011), Introduction to generalized topological space, Applied General Topology, Vol-12(1), (pp.49-66).
- [22]. ShankarR.(2024), Remarks on strong $\pi^* - \mathcal{H}$ -open sets in GTS via Hereditary class,International journal of scientific research in mathematical science and statistical science, Vol-11(2), (pp. 27-31),
- [23]. KimY.K., and KimW.K. (2012), On operations induced by hereditary classes on generalized topological space, Acta Math. Hungar, Vol-137(1-2), (pp. 130-138).

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website

www.ijeast.com



INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

✉ editor@ijeast.com

🌐 www.ijeast.com

📍 India



2455-2143