



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 11 ISSUE : 01 Print / Issue Publication Date: 02-Jun-2026



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2026.v11i01.022

Indexed In



WWW.IJEAST.COM

editor@ijeast.com

THE PHYSICS OF ANTIMATTER ANNIHILATION AND ENERGY RELEASE

Dharmendra Kumar
Research Scholar,
CSIR Net JRF December 2024

Abstract: In this study, the physics of matter – antimatter annihilation as well as the tremendous amount of energy produced when matter – antimatter particles interact, is explored through the lens of relativistic physics, conservation laws, and quantum electrodynamics. According to Einstein's equation, $E=mc^2$, the analysis shows that almost all of the rest mass is turned into energy when annihilation takes place. In the annihilation of 1 g of matter with 1 g of antimatter, it is estimated that about 1.8×10^{14} J (several kilotons of TNT) of energy is released. The two 511 keV γ photons are the result of annihilation of an electron with a positron, and converses momentum and charge. The relativistic modeling suggests that the energy of particles rapidly rises for relativistic momenta above 10^{-21} kg·m/s. The results of the gamma-ray attenuation analysis show that when the attenuation coefficient μ is 0.1, high density shielding materials can reduce the intensity of the radiation below 10% after 20cm thickness. Further calculations using the thermal deposition show that a material with a mass of 0.1 g can be heated to almost 1000 K when it absorbs 2000 J of energy. The study underscores the unparalleled energy density and radiation properties of anti-matter annihilation in particle physics, astrophysics, medicine, and the development of future energy technologies.

Keywords: Antimatter physics, Matter–antimatter annihilation, Mass–energy equivalence, Gamma-ray radiation, Quantum electrodynamics, Relativistic energy release

I. INTRODUCTION

The subject of annihilation of antimatter is one of the most interesting and important in contemporary physics. Antimatter, first theorized in 1928 by Paul Dirac, is particles that have the same mass as particles of matter, but the opposite electric charge and quantum properties. There is an anti-particle for each fundamental particle of the universe [2]. The positron is the antiparticle of the electron, for instance, and the antiproton is the antiparticle of the proton. If matter and antimatter meet, the masses of both are converted into huge amounts of energy, a phenomenon known as annihilation, which is described by the Mass–

energy equivalence postulated by Albert Einstein [3]. This is described by the well known formula $E = mc^2$.

Here, E is energy, m is mass, c is the speed of light. This equation shows that a very small amount of mass can be transformed to a great amount of energy [4].

Nearly 100% of the mass in the antimatter annihilation process can be converted to energy, which is one of the most efficient energy conversion processes known in physics [5]. A high-energy photon or gamma ray would be just as striking a sign of annihilation as would be high-energy subatomic particles. In conventional chemical reactions, only a small amount of matter is converted into useful energy; in annihilation, energy is released in the form of high-energy photons, gamma rays, and other subatomic particles [6]. The incredible energy density of the antimatter has been the focus of research in particle physics, astrophysics, cosmology and new energy systems, among others [7]. Antimatter, they believe, may hold a key to the answers to some of the universe's deepest mysteries, such as the imbalance of matter and antimatter that occurred after the Big Bang: [8]

The phenomenon of antimatter annihilation has been observed in laboratory experiments throughout the world and is not theoretical. Groups like CERN have been able to create and experiment with antiparticles under controlled conditions [9]. Experiments with electron–positron collisions have proven that annihilation emits powerful showers of gamma radiation and secondary particles. This has significantly increased the knowledge of quantum physics, particle interactions and high energy physics [10]. In addition to that, the detection and capture of antihydrogen atoms has given rise to new avenues to explore basic symmetries of nature, and to investigate the laws of the universe.

Astrophysicists use the concept of antimatter annihilation to help understand some of the cosmic phenomena. The formation of high-energy gamma rays observed in space are frequently linked with the interactions of matter–antimatter near black holes, neutron stars and other energetic astronomical objects [11]. Others scientists try to determine if there's any other place in the universe where antimatter exists. It is thought to have had an effect on the early evolution of the cosmos shortly after the Big Bang, when an almost equal amount of matter and antimatter were created [12]. The question as to why matter prevails in the current



universe, however, has not been solved yet and is one of the biggest mysteries in cosmology.

In addition to being an interesting and important scientific field, antimatter annihilation also has practical implications in today's technology and medicine. A prominent application is Positron Emission Tomography (PET) which is a medical imaging technique for recognizing disease and checking biological processes in the human body [13, 14]. The potential of energy generation and propulsion of deep space missions with antimatter is also being investigated because of its high energy output. Even with these promising applications, the production and storage of antimatter is still very difficult and costly, as antimatter is destroyed on contact with the normal matter. The aims of this research paper are as follows:

- To study the fundamental concepts, properties, and behavior of antimatter and antiparticles in modern physics.
- To understand the process of matter–antimatter annihilation and analyze the mechanism of energy release based on mass–energy equivalence.
- To examine the significance of antimatter annihilation in particle physics, astrophysics, cosmology, and experimental research.
- To explore the practical applications, challenges, and future potential of antimatter in medicine, energy production, and space technology.

II. LITERATURE REVIEW

This literature review reveals the importance of the contributions made in recent years to the theoretical groundwork, the experimental investigation and the utilization of antimatter physics and annihilation and energy release. Victor Flambaum et al. (2021) [15] have studied the creation of positron clouds in the quark nugget model of dark matter and find that the electrons' annihilations with positrons in quark nuggets cannot provide the whole picture of the observed 511 keV photon emissions in the galaxy. The alternative annihilation processes proton capture and pion decay were also stressed by their work. This work Tanu Zhang et al., 2022 [16] has provided a detailed overview of the discovery, production, storage and future applications of antimatter, including its use as a source of energy. In the same manner, the properties of antiparticles, the production and storage of them, annihilation of particles and antiparticles, and their technological applications were discussed by Hamouda et al. (2020) [17]. Youvan et al. (2024) [18] discussed the tremendous amount of energy that antimatter possesses, noting that it would take only one gram of antimatter to yield the energy that's comparable to several kilotons of TNT! They also explored the difficulties associated with the production, storage and potential applications of antimatter in space propulsion, medicine and energy systems. In addition, Hossain et al., 2023 [19]

discussed the interconnections among antimatter, dark matter and dark energy and their relevance in comprehending the structure and evolution of the universe. Furthermore, Gregori et al. 2022 [20] studied the annihilation of particles and antiparticles in a more general theoretical and cosmological setting and correlate the annihilation processes with universal dynamics and energy conversions.

New experiments and cutting-edge uses of antimatter annihilation have also been the subject of recent research. An alternative to the dark matter/dark energy theories was proposed by Gouveia et al. (2025) [21] with their wave-based cosmological model that accounts for galactic rotation and cosmic expansion by the release of energy from the annihilation of primordial matter–antimatter. Xu et al. (2024) optimized the $\beta^+\text{-}\gamma$ coincidence positron annihilation lifetime spectroscopy system for large coincidence counting rate and high time resolution, which is beneficial to enhancing the study on defect characterization in material science. Experiments with antiproton–nucleus annihilation at rest with slow antiprotons from CERN were performed by Amsler et al. [23] and were compared to Geant4 and FLUKA simulations, where they found that the models require more accurate modelling of annihilation in the low-energy regime. Likewise, Ferreira et al. (2025) [24] proposed a deep learning-based annihilation vertex reconstruction system for the ALPHA-g experiment at CERN, showing the system can achieve a higher accuracy in determining the location of annihilation of antihydrogen. As all the above studies show, the field of antimatter annihilation is an exciting one which has many implications for the field of particle physics, cosmology, medical science and future energy technologies.

Although significant progress has been made in the study of antimatter, there are still many aspects of the full mechanism of the annihilation of matter and antimatter and the resulting release of energy that remain unknown. Previous work dedicated to the theoretical concepts and/or experimental detection of antimatter, with few studies that consider efficient production of antimatter, safe storage, and practical energy use. Even current simulation models are not successful in explaining the annihilation processes at low energy. Thus, additional research efforts are required to create better theoretical models, as well as advanced technologies, for harnessing antimatter energy for scientific and technological use.

III. RESEARCH METHODOLOGY

This research is based on the theories of physics, relativistic quantum mechanics, and conservation-law analysis to examine the annihilation of matter and antimatter and the release of energy. To analyze annihilation reactions, photon emission, the relativistic behaviour of particles, the attenuation of gamma-rays or the transfer of thermal energy, during high-energy interactions with antimatter,



mathematical modeling, quantum electrodynamics and numerical estimation methods are applied.

$$Q_i = Q_f \tag{6}$$

3.1 Research Design

This study delves into the theoretical and theoretical physics aspects of annihilation of matter and antimatter in light of conservation laws and the energy that is released. The modelling of annihilation reactions, photon emission, radiation behaviour and energy conversion efficiency is carried out with the help of analytical derivations and numerical estimation methods. In addition the method is tested for relativistic particle interactions and the dynamics of energy transfer for the gamma ray during annihilation.

- **Photon Energy Distribution**

During electron–positron annihilation at rest, two gamma rays are simultaneously created each moving in one direction. Each photon has the same energy as an electron's rest-mass energy. The photons emitted also have momentum and thus conserves momentum during annihilation. This energy distribution is necessary for the gamma-ray spectroscopy and high-energy particle interaction analysis.

$$E_\gamma = m_e c^2 \tag{7}$$

3.2 Theoretical Framework

- **Matter–Antimatter Interaction**

Antimatter particles have the same mass as the particle of matter but have the opposite quantum numbers and electric charge. A particle and its antiparticle when they collide are annihilated and their mass is turned into high-energy photons and other particles. The characteristic feature of electron–positron annihilation is the production of two gamma-ray photons, which, in accordance with the conservation of energy and momentum, share its energy and momentum equally.

- **Relativistic Energy Relations**

Antiparticles initially have a relativistic energy that will be obtained from the relativistic energy–momentum relation. This equation is the sum of kinetic energy, momentum and rest-mass energy equations and is used to describe the behavior of high-energy particles with great accuracy. It has many applications in particle physics, such as calculating annihilation products, collision dynamics, and the energy of the emitted photons or secondary particles.

$$E^2 = p^2 c^2 + m^2 c^4 \tag{8}$$



- **Mass–Energy Equivalence**

The energy release in the process where matter and anti-matter collide each other is calculated from the mass-energy relation discovered by Einstein. The annihilation is very efficient because the total rest mass of both the particles are converted into energy directly. The energy emitted for the same amount of matter and antimatter is proportional to the speed of light to the power of 2 and the two masses combined.

3.4 Quantum Electrodynamics (QED) Modeling

- **Interaction Cross-Section**

In the context of Quantum electrodynamics, the “interaction cross section” is a measure of the likelihood of a collision where an electron and a positron annihilate. The annihilation rate is proportional to the classical electron radius and the relative velocity of the particles in interactions of low energy. The prediction of collision frequency, annihilation probability and gamma-ray production in accelerator and antimatter experiments is crucial and requires a cross-section analysis.

$$E = mc^2 \tag{2}$$

$$E_{\text{total}} = (m_m + m_a)c^2 \tag{3}$$

$$\sigma = \pi r_e^2 \left(\frac{1-v^2/c^2}{v/c} \right) \tag{9}$$

3.3 Mathematical Modeling

- **Conservation Laws**

The fundamental conservation laws that apply to matter – antimatter annihilation include conservation of energy, conservation of momentum, and conservation of electric charge. These laws apply to the properties of the emitted photons and secondary particles in annihilation reactions. The amount of total energy, momentum and charge before and after interaction are conserved, so that particle transformations are physically valid as well as the modeling of relativistic interactions.

- **Feynman Diagram Analysis**

The Feynman diagram analysis approach explains the interaction of antimatter via quantum electrodynamics and virtual photon exchange. The particles can create a virtual photon as an intermediate stage in the annihilation process and then the virtual photon can create higher-mass particle pairs. Transition amplitudes, interaction probabilities, and scattering behaviors are computed in high energy particle collision processes by means of the perturbation theory. The perturbation theory is applied to calculate the transition amplitude, the interaction probability, and the scattering behavior in the high energy particle collision process.

$$E_i = E_f \tag{4}$$

$$\vec{p}_i = \vec{p}_f \tag{5}$$



3.5 Energy Release Estimation

The energies involved with the annihilation of matter and antimatter are determined by the mass-energy equivalence principle. In the case of annihilation, 1 gram of matter and 1 gram of antimatter turn into almost 2 grams of energy; the resulting energy output is vast, much greater than is released in regular nuclear reactions. Theoretically, it is 100% efficient at converting energy as almost all the rest mass is converted to radiation energy.

$$\eta = \frac{E_{\text{useful}}}{E_{\text{total}}} \times 100\% \tag{10}$$

3.6 Radiation Transport Analysis

Radiation transport analysis (RTA) is used to analyze how gamma rays produced by the annihilating of the antimatter travel and are affected by material near the annihilation point. As the radiation passes through the matter, the intensity of the gamma rays decreases in an exponential manner depending of the attenuation properties of the material and the thickness of the material. The energy absorbed also generates thermal heating effects, which raises the temperature of the material and plays a role in the formation of the plasma in high-energy environments.

$$Q = mc\Delta T \tag{11}$$

IV. RESULT AND DISCUSSION

Figure 2 illustrates the relationship between mass and released energy based on Einstein’s mass-energy equivalence equation, $E = mc^2$. The logarithmic trend shows that energy increases proportionally with mass. For example, a mass of 10^{-6} kg produces approximately 10^{11} J of energy, while 10^2 kg generates nearly 10^{19} J. The graph demonstrates the enormous energy potential stored within matter and highlights why antimatter annihilation is considered one of the most powerful theoretical energy-generation processes in modern physics.




Figure 2: Mass-Energy Equivalence Graph

Figure 3 compares relativistic total energy with constant rest energy for an electron as momentum increases. The red line represents rest energy, remaining nearly constant at approximately 8.2×10^{-14} J, while the blue dashed curve shows relativistic energy increasing rapidly with momentum according to $E^2 = p^2c^2 + m^2c^4$. At low momentum (10^{-25} kg·m/s), relativistic energy is close to rest energy, but near 10^{-16} kg·m/s, energy rises above 10^{-8} J, demonstrating strong relativistic effects at high particle velocities.




Figure 3: Relativistic Energy-Momentum Relation

Figure 4 illustrates the relationship between annihilated mass and total energy released during matter-antimatter annihilation using $E_{\text{total}} = (m_m + m_a)c^2$. Energy increases linearly with mass on a logarithmic scale. An annihilated mass of 10^{-9} g produces nearly 10^6 J, while 1 g releases approximately 10^{20} J of energy. The graph demonstrates the extremely high energy density of antimatter reactions, greatly exceeding conventional chemical and nuclear energy sources.




Figure 4: Energy Released from Matter-Antimatter Annihilation

Figure 5 shows the exponential reduction of normalized gamma-ray intensity as material thickness increases, based on the attenuation equation $I = I_0 e^{-\mu x}$. Higher attenuation coefficients cause faster intensity decay. For $\mu = 0.20 \text{ cm}^{-1}$, intensity decreases below 10^{-4} after 20 cm, while for $\mu = 0.05 \text{ cm}^{-1}$, intensity remains near 10^{-3} . The results demonstrate how dense shielding materials absorb annihilation-generated gamma radiation more effectively, reducing radiation penetration and improving protection against high-energy photons.




Figure 5: Gamma-Ray Attenuation in Matter

Figure 6 illustrates the relationship between absorbed energy and temperature increase for different material masses using the heat transfer equation $Q = mc\Delta T$. Temperature rise increases linearly with absorbed energy, while smaller masses experience greater heating effects. For an absorbed energy of 2000 J, a 0.1 g material reaches nearly 1000 K, whereas a 1 g material increases to approximately 400 K. The results demonstrate how annihilation-generated radiation can produce intense thermal and plasma effects in low-mass materials.




Figure 6: Thermal Energy Deposition and Temperature Rise

V. CONCLUSION

The study ends by showing that matter-antimatter annihilation is among the most efficient ways of converting

energy to energy in contemporary physics. By converting almost 1.8×10^{14} J of energy into pure annihilation and converting nearly 100% of the matter and antimatter, Einstein's mass-energy equivalence relation makes it clear that annihilation of 1 g matter with 1 g antimatter can produce almost 1.8×10^{14} J of energy. Conservation principles and predictions of quantum electrodynamics are confirmed when an electron-positron pair annihilates, creating a pair of oppositely traveling γ photons, each with 511 keV in energy. According to the relativistic energy analysis, the energy increases rapidly with high particle momentum, and according to gamma-ray attenuation results, the penetration of radiation is significantly suppressed below 10% for a dense shielding of 20 cm thickness. Thermal energy-deposition modelling shows that strong absorption of radiation can lead to temperature rises of up to 1000K in low mass materials. The discovery highlights the vast applications of antimatter in the fields of advanced energy systems, medical imaging, astrophysical research, and future space propulsion, yet some obstacles in antimatter production, storage, and radiation risks of handling remain impractical at present.

VI. REFERENCES

- [1]. Kragh, Helge. "Antimatter in astronomy and cosmology: the early history." *Annals of Science* 83, no. 2 (2026): 399-423.
- [2]. Baker, David John, and Hans Halvorson. "Antimatter." *The British Journal for the Philosophy of Science* 61, no. 1 (2010): 93-121.
- [3]. Salah, Ahmed Ragab Mahmoud. "E= mc² Explained: Einstein's Mass-Energy Equivalence and Its Revolutionary Role in Physics, Nuclear Power, and the Universe." (2025).
- [4]. Spence, John CH. "Speed limit: How the search for an absolute frame of reference in the Universe led to Einstein's equation 'E= mc²'-a history of measurements of the speed of light." In *Journal and Proceedings of the Royal Society of New South Wales*, vol. 152, no. 473/474, pp. 216-241. 2019.
- [5]. Deutsch, Claude, and Naeem A. Tahir. "Fusion reactions and matter-antimatter annihilation for space propulsion." *Laser and Particle Beams* 24, no. 4 (2006): 605-616.
- [6]. Karmaker, Nanda, Kazi M. Maraz, Farhana Islam, Md Marjanul Haque, Md Razzak, M. Mollah, M. R. I. Faruque, and Ruhul A. Khan. "Fundamental characteristics and application of radiation." *GSC Advanced Research and Reviews* 7, no. 1 (2021): 064-072.
- [7]. Yang, Xiaodong, Yuchen Yang, Zhen Luo, and Yuanbo Bi. "Research Again Origin of the Asymmetry Between Matter and Antimatter—Energy Basic State Field of the Universe (II)." (2025).



- Reports in Advances of Physical Sciences 7 (2023): 2330001.
- [8]. Canetti, Laurent, Marco Drewes, and Mikhail Shaposhnikov. "Matter and Antimatter in the Universe." *New Journal of Physics* 14, no. 9 (2012): 095012.
- [9]. Gourlay, Stephen, Tor Raubenheimer, and Vladimir Shiltsev. "Challenges of future accelerators for particle physics research." *Frontiers in Physics* 10 (2022): 920520.
- [10]. Feldman, Gary J., and Martin L. Perl. "Electron-positron annihilation above 2 GeV and the new particles." *Physics Reports* 19, no. 5 (1975): 233-293.
- [11]. Canetti, Laurent, Marco Drewes, and Mikhail Shaposhnikov. "Matter and Antimatter in the Universe." *New Journal of Physics* 14, no. 9 (2012): 095012.
- [12]. Kurki-Suonio, Hannu, and Elina Sihvola. "Antimatter regions in the early universe and big bang nucleosynthesis." *Physical Review D* 62, no. 10 (2000): 103508.
- [13]. Sunderland, John. "Medical imaging with antimatter." *Physics today* 78, no. 9 (2025): 28-35.
- [14]. Eissa, Mostafa Essam. "Antimatter: The potential impact on the future of medical and pharmaceutical industries." *Universal Journal of Pharmaceutical Research* (2025).
- [15]. Flambaum, V. V., and I. B. Samsonov. "Radiation from matter-antimatter annihilation in the quark nugget model of dark matter." *Physical Review D* 104, no. 6 (2021): 063042.
- [16]. Zhang, Ruichen. "Antimatter and Its Application-Collecting Antimatter and Storage It as Energy Source." In *Journal of Physics: Conference Series*, vol. 2386, no. 1, p. 012074. IOP Publishing, 2022.
- [17]. Hamouda, Samir A. "Matter-antimatter and its Potential Applications." Department of Physics, University of Benghazi, Libya (2020).
- [18]. Youvan, Douglas C. "Pathways to Antimatter Abundance: Exploring Breakthroughs for Large-Scale Antimatter Production, Applications, and Explosive Potential." (2024).
- [19]. Hossain, Khandakar Akhter. "Evaluate the Mystery of Creation of Universe and Existence of Antimatter Dark Matter and Dark Energy." *International Journal of Current Science Research and Review* 6, no. 06 (2023).
- [20]. Gregori, Giovanni Pietro, Bruce Allen Leybourne, Willie Soon, and Valentino Straser. "The foundations of physics and axiomatics III-Superluminal phenomena, mechanisms, matter-antimatter, cosmological implications." *New Concepts in Global Tectonics Journal* 10, no. 3 (2022).
- [21]. Gouveia, Gui Furne. "The Role of Energy Density Diffusion in Galactic Dynamics and Cosmic Expansion: A Unified Theory for MOND and Dark Energy." (2025).
- [22]. Xu, W., Y. H. Li, Z. W. Pan, J. D. Liu, H. J. Zhang, and B. J. Ye. "Optimization study of a new $\beta^+-\gamma$ coincidence positron annihilation lifetime spectrometer using Geant4." *Journal of Instrumentation* 19, no. 06 (2024): P06044.
- [23]. Amsler, Claude, Horst Breuker, Marcus Bumbar, Matti Cerwenka, Giovanni Costantini, Rafael Ferragut, Markus Fleck et al. "Antiproton annihilation at rest in thin solid targets and comparison with Monte Carlo simulations." *The European Physical Journal A* 60, no. 11 (2024): 225.
- [24]. Ferreira, Ashley, Mahip Singh, Yukiya Saito, Andrea Capra, Ina Carli, Daniel Duque Quiceno, W. Fedorko et al. "Antimatter annihilation vertex reconstruction with deep learning for ALPHA-g radial Time Projection Chamber." *Journal of High Energy Physics* 2025, no. 7 (2025): 1-16.

IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

ABOUT IJEAST

International Journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high-quality research papers in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

FOCUS AREAS

- Engineering
- Applied Science
- Technology
- Innovation & Development
- Interdisciplinary Studies



PEER REVIEWED

All submissions are rigorously peer reviewed to ensure quality.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website
www.ijeast.com



INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY

✉ editor@ijeast.com

🌐 www.ijeast.com

📍 India



2455-2143