



IJEAST

INTERNATIONAL JOURNAL
OF ENGINEERING APPLIED SCIENCE
AND TECHNOLOGY



VOLUME : 7 ISSUE : 03 Print / Issue Publication Date: 29-Aug-2022



ISSN : 2455-2143



DOI : 10.33564/IJEAST.2022.v07i03.018

Indexed In



WWW.IJEAST.COM

editor@ijeast.com



ON NON-HOMOGENEOUS TERNARY CUBIC DIOPHANTINE EQUATION

$$w^2 - z^2 + 2wx - 2zx = x^3$$

S.Vidhyalakshmi
 Assistant Professor,
 Department of Mathematics
 Shrimati Indira Gandhi College,
 Affiliated to Bharathidasan University, Trichy-620 002,
 Tamil Nadu, India.

M.A.GOPALAN
 Professor,
 Department of Mathematics,
 Shrimati Indira Gandhi College,
 Affiliated to Bharathidasan University, Trichy-620 002,
 Tamil Nadu, India.

J. Shanthi, T. Mahalakshmi
 Assistant Professor,
 Department of Mathematics,
 Shrimati Indira Gandhi College,
 Affiliated to Bharathidasan University, Trichy-620 002,
 Tamil Nadu, India.

Abstract— The non-homogeneous ternary cubic Diophantine equation $w^2 - z^2 + 2wx - 2zx = x^3$ is analyzed for its patterns of non-zero distinct integral solutions. A few relations between the solutions and special number patterns are presented.

Keywords— Ternary cubic Non- Homogeneous cubic, Integral solutions.

Notations:

$$t_{m,n} = n(1 + \frac{(n-1)(m-2)}{2})$$

$$p_n^r = \frac{n(n+1)(n(r-2) - (r-5))}{6}$$

$$CP_k^4 = \frac{4k^3 + 2k}{6}, \quad CP_k^8 = \frac{8k^3 - 2k}{6}$$

I. INTRODUCTION

The Diophantine equation offers an unlimited field for research due to their variety [1-3]. In particular, one may refer [4-15] for cubic equations with three unknowns. This

communication concerns with yet another interesting equation $w^2 - z^2 + 2wx - 2zx = x^3$ representing non-homogeneous cubic with three unknowns for determining its infinitely many non-zero integral points. A few relations between the solutions and special number patterns are presented.

II. METHOD OF ANALYSIS

The given non-homogeneous ternary cubic diophantine equation is

$$w^2 - z^2 + 2wx - 2zx = x^3 \quad (1)$$

On completing the squares, (1) is written as

$$P^2 - Q^2 = x^3 \quad (2)$$

Where

$$P = w + x, Q = z + x \quad (3)$$

Write (2) as the system of double equations as below:

$$P + Q = x^3,$$

$$P - Q = 1$$

Solving the above system of equations, one obtains

$$x = 2k + 1 \quad (4)$$

and

$$P = 4k^3 + 6k^2 + 3k + 1, Q = 4k^3 + 6k^2 + 3k \quad (5)$$



In view of (3), we have

$$w = 4k^3 + 6k^2 + k, z = 4k^3 + 6k^2 + k - 1 \quad (6)$$

Thus, (4) and (6) represent the integer solutions to (1).

Relations between the solutions and special number patterns:

I. Each of the following expressions is a square multiple of 2

$$(i). w - x * t_{3,2k}$$

$$(ii). z + 1 - x * t_{3,2k}$$

$$(iii). w + t_{3,2k} - 3 * P_{2k}^3$$

$$(II). w = -12t_{3,k} + 24 * P_k^3 - k$$

$$(III). w = -t_{14,k} + 24 * P_k^3 - 12k$$

$$(IV). w - x = t_{6,k} + 8 * P_k^5 - 1$$

(V). Each of the following is a square multiple of 6

$$(i). w - 6CP_k^4 + k$$

$$(ii). w + z + x - 6CP_k^8 + 12t_{3,k}$$

For simplicity and brevity, the other choices of solutions to (1) are exhibited below:

Choice 1:

$$x = k, z = t_{3,k-1} - k, w = t_{3,k} - k$$

Choice II:

$$x = 2na, z = 2n^3a^2 - (2n+1)a, w = 2n^3a^2 - (2n-1)a$$

Choice III

$$x = 2na, z = a^2 - 2(n^3 + n)a, w = a^2 + 2(n^3 - n)a$$

Choice IV:

$$x = 2na, z = 2n^3a^3 - (2na+1), w = 2n^3a^3 - (2na-1)$$

Choice V:

$$x = 2na, z = a^3 - 2n(n^2 + a), w = a^3 + 2n(n^2 - a)$$

III. CONCLUSION

In this paper, we have made an attempt to obtain all integer solutions to (1). To conclude, one may search for integer solutions to other choices of ternary cubic diophantine equations.

IV. REFERENCE

- [1]. Dickson, L.E.; (1952) History of Theory of Numbers, vol 2, Chelsea publishing company, New York.
- [2]. Mordell, L.J.; (1969) Diophantine Equations, Academic press, London..
- [3]. Carmichael, R.D.; (1959), The theory of numbers and Diophantine analysis, New York, Dover.
- [4]. Gopalan, M.A.; Srividhya, G. (2011), Integral solutions of ternary cubic diophantine equation $x^3 + y^3 = z^2$ Acta Ciencia Indica, Vol XXXVII, No.4, (Pp 805-808).
- [5]. Gopalan, M.A.; Vidhyalakshmi, S.; Thiruniraiselvi, N. (2013), On Non-Homogeneous Cubic Equation With Three Unknowns $x^2 - y^2 + z^2 = 2kxyz$, BOMSR, Vol.1(1), (Pp 13-15).
- [6]. Gopalan, M. A; Vidhyalakshmi, S; Lakshmi, K, (2013), Latic points on the non-homogeneous cubic equation $x^3 + y^3 + z^3 + (x + y + z) = 0$, Impact.J.Sci. Tech, Vol.7, No. 1, (Pp 21-25).
- [7]. Gopalan, M. A; Vidhyalakshmi, S; Lakshmi, K, (2013), Latic points on the non-homogeneous cubic equation $x^3 + y^3 + z^3 - (x + y + z) = 0$, Impact.J.Sci. Tech, Vol.7, No. 1, (Pp 51-55).
- [8]. Vidhyalakshmi, S; Usharani, T.R; Gopalan, M.A, (2014), Integral Solutions Of The Ternary Cubic Equation $5(x^2 + y^2) - 9xy + x + y + 1 = 35z^3$, IJRET, Vol:03, Issue:11, (Pp 449-452).
- [9]. Gopalan, M.A; Thiruniraiselvi, N; Sridevi, R, (2015), On The Ternary Cubic Equation $5(x^2 + y^2) - 8xy = 74(k^2 + s^2)z^3$, IJMRE, Vol.1(1), (Pp 317-319).
- [10]. Gopalan, M. A; Vidhyalakshmi, S; Shanthi, J; Maheswari, J, (2015), On ternary cubic Diophantine equation $3(x^2 + y^2) - 5xy + x + y + 1 = 12z^3$, IJAR, Volume 1, Issue 8, (Pp 209-212).
- [11]. Janaki, G and Saranya, P; (2016), On the Ternary Cubic Diophantine Equation $5(x^2 + y^2) - 6xy + 4(x + y) + 4 = 40z^3$, International Journal of Science and Research- online, Vol 5, Issue3, (Pp 227-229).
- [12]. Gopalan, M.A, Sharadha Kumar, (2018), "On the non-homogeneous ternary cubic equation $3(x^2 + y^2) - 5xy + x + y + 1 = 111z^3$ International Journal of Engineering and Techniques, 4(5), (Pp 105-107).
- [13]. Sharadha Kumar, Gopalan, M.A; (2019), "On The Cubic Equation $x^3 + y^3 + 6(x + y)z^2 = 4w^3$ ", JETIR, 6(1), (Pp 658-660).
- [14]. Vijayasankar, A; Dhanalakshmi, G; Sharadha Kumar, Gopalan, M.A; (2020), On The Integral Solutions To The Cubic Equation With Four Unknowns $x^3 + y^3 + (x + y)(x - y)^2 = 16zw^2$ International Journal For Innovative Research In Multidisciplinary Field, 6(5), (Pp 337-345).
- [15]. Vijayasankar, A; Sharadha Kumar, Gopalan, M.A, (2021), "On Non-Homogeneous Ternary Cubic Equation $x^3 + y^3 + x + y = 2z(2z^2 - a^2 + 1)$ International Journal of Research Publication and Review 2(8), (Pp 592-598).

IJEAST

INTERNATIONAL JOURNAL OF ENGINEERING APPLIED SCIENCE AND TECHNOLOGY



AUTHORS

 S.Vidhyalakshmi

 M.A.GOPALAN

 J. Shanthi

 T. Mahalakshmi



ABOUT IJEAST

International journal of Engineering Applied Science and Technology (IJEAST) is a peer-reviewed, open access journal that publishes high - quality research paper in the field of Engineering, Applied Science and Technology.

IJEAST aims to provide a platform for researchers, academicians, and professionals to share their innovative ideas, research findings, and practical experiences with the global scientific community.

JOURNAL METRICS

H

H-INDEX

33

i10

i10-INDEX

154

“

IJEAST
CITATIONS

9,647

(IJEAST CITATIONS)



PEER-REVIEWED
& OPEN ACCESS

PUBLISHED
MONTHLY



PEER REVIEWED

All submission are rigorously peer reviewed to ensure quality.



AUTHORS ARE OUR PRIORITY

We value our authors and recognize their contribution to the advancement of research and knowledge.



OPEN ACCESS

Free and unrestricted access to research for all.



GLOBAL REACH

Connecting researchers and professionals worldwide.



TIMELY PUBLICATION

We ensure a swift and efficient publication process.



For more information, visit our website
www.ijeast.com



editor@ijeast.com



www.ijeast.com



India



2455-2143