



DENTAL DISEASES DETECTION SYSTEM USING pH SALIVA - A REVIEW

Panchashila P.Wakode

Department of electronics and telecommunication
Maharashtra institute of technology, Aurangabad,
Maharashtra, India

Shilpa kodgire

Department of electronics and telecommunication
Maharashtra institute of technology, Aurangabad,
Maharashtra, India

Abstract— Saliva, science has revealed, is a great deal greater than water. A Saliva is the complicated mixture of fluids that surrounds the oral tissues, and it originates from essential and minor salivary glands and non glandular sources such as crevicular fluids, oral microorganisms and host cells. The consistency of saliva can be watery, thick, sticky or frothy depending on its composition. the amount of proteins in saliva essentially will determine its thickness or frothiness. A basal unstimulated secretion is produced constantly to moisturize and lubricate the oral tissues for greater than 90 percentage of the day. gustatory, olfactory or pharmacological stimuli enlarge the production and secretion of saliva. Saliva is an increasing number of extra known as an captivating diagnostic fluid. The presence of a wide variety of sickness signaling salivary biomarkers that precisely replicate everyday and ailment states in people and the sampling advantages in contrast to blood sampling are some of the motives for this recognition. This explains the burgeoning research difficulty in assay features and technological developments for the detection of quite a variety salivary biomarkers to enhance scientific diagnosis, management, and treatment. Saliva is an amorphous fluid and, we comprehend two about its plenty of properties, rather little is known about the element of its molecular content. This complete know-how made it tough or even not possible to behavior scientific discovery research to stumble on salivary biomarkers that need to be used for stickiness detection With funding assist from NIDCR, numerous research centers in the United States tackled this assignment by way of first defining and cataloging all the proteins in saliva. It is packed with proteins that help manage the teeming hordes of microbes in our mouths. It is stuffed with materials that make us spit gristly, end our enamel from dissolving and assist heal wounds. It is packed with a plethora of hormones and other chemicals revealing something from whether one smokes to whether one is stressed. Saliva is a clear, slightly acidic mucoserous exocrine Biofluid produced in the oral cavity by means of three predominant (parotid, submandibular and sublingual).It is a complicated fluid containing an complete library of hormones, proteins,

enzymes, antibodies, antimicrobial constituents, and cytokines.

Keywords— Human saliva, arduino, pH probe electrode

I. INTRODUCTION

Saliva is biologic fluid in the oral cavity. composed of a mixture of secretory products from the major and minor salivary glands. Saliva plays key roles in lubrication, mastication. taste perception. Prevention of oral infection of dental caries. Saliva has many important roles in oral health like lubrication, physical protection, cleansing, buffering, tooth integrity, maintenance of pH and antibacterial action [2].Several authors have mentioned significant correlations between the composition of certain substances in saliva and blood. Saliva seems to be a promising diagnostic fluid in the monitoring of general public health, early disorder detection, bio monitoring of pesticides, monitoring of drugs and smoking and various forensic and scientific functions[1].Dental caries is the most popular dental disease affecting human race even though the incidence of dental caries has extensively reduced, it is nevertheless a major problem[3]. Normal healthy men and ladies between 18– 40 years who had been not below medications that could affect salivation had been selected. Pregnant ladies, smokers or betel leaf chewers have been excluded. The secretion fee of stimulated saliva is expressed as milliliter per min. Stimulated salivary secretion rate higher than 1 ml / min was considered to be normal. Human saliva is no longer simply the fluid in our mouth, but it mirrors our body's fitness and well being. Bio molecules that are circulating in the blood are also observed in human saliva. It consists of approximately about 2,000 proteins, and most significantly, 26% of these proteins are also found in blood, therefore emphasizes the saliva's significance as an brought organic useful resource for disease analysis and monitoring, as nicely as an remaining diagnostic medium to set up a person's response to treatment. The subject of saliva diagnostics (SDs) started out in the early 60s when salivary calcium stages had been determined to be increased in cystic fibrosis patients, and 50 years on now how the field has



unmitigated to unpredicted distance due to the development of increasingly sensitive detection techniques[7].

The significance of Saliva to be used as a diagnostic fluid:

Saliva as a diagnostic fluid presents a unique advantages over serum due to the fact it can be accrued non-invasively and does not require exclusive gear for series and storage as not like blood, saliva does not clot. Advantageous for human beings in whom blood drawing is challenging as in obese, hemophiliacs and sufferers who are worried of prick. WS can be used for analysis of systemic diseases, because it consists of serum constituents, for some diagnostic purposes, salivary biomarkers proved more useful than serum analysis. These components are derived from the local vasculature of the salivary glands and gingival crevicular fluid.8 Many advantages of saliva as a clinical tool over serum and tissues are a noninvasive collection of sample, smaller sample aliquots, good cooperation with patients, fee effectiveness, easy storage and transportation, repeated sampling for monitoring over time, greater sensitivity, and correlation with ranges in blood. Most frequently referred to criticism of using saliva as a diagnostic fluid is these biomarkers are present in quantities that are too low to be detected reliably. However, this is no longer a problem due to the improvement of an increasing number of sensitive detection techniques. Hence today in the technology of nanotechnology and genomics, field of SDs is promising a dramatic trade in ailment prognosis and clinical monitoring. With cutting-edge research, the gap is closing unexpectedly between the use of saliva and other biofluid for disease diagnostics. The rising discipline of microbiology and nano technology based totally biosensor will overcome the detection barriers. Two prerequisites exist before the intention of SDs can be achieved: identification of specific biomarkers related with a health or sickness state and the development of technologies that can discriminate between the bio makers. A current initiative of the National Institute of Dental and Craniofacial Research (NIDCR) has created a roadmap to gain these desires via the use of oral fluids as the diagnostic medium to scrutinize the health and/or ailment reputation of patients. This is an best probability to optimize today's saliva-based biosensors for salivary biomarkers that discriminate between diseases. This is an interesting time, as we are seeing the functions of SDs for oral diseases, which will be followed soon via the application to high-impact systemic diseases, using relatively informative panels of salivary proteomic and genomic biomarkers. This will enable researchers to bridge oral fitness research with systemic disorder diagnostics by way of a biofluid that filters, techniques and secretes itself from the vasculature that nourishes the salivary glands into the oral cavity.[7]

II. LITERATURE REVIEW

Khushbu Yadav KMTRC, Janakpurdham, Nepal, Satyam Prakash-Dental caries, a persistent malady is interesting among human and is one of the foremost common critical global oral well being issues within the world nowadays. It is the devastation of dental difficult a cellular tissue by acidic by-products from the bacterial maturation of dietary carbohydrates particularly sucrose. It advances gradually in most of the people which comes about from an environmental lop sidedness within the balance between tooth minerals and verbal biofilms which is characterized by microbial movement, coming about in variances in plaque pH due to bacterial corrosive generation, buffering activity from spit and the encompassing tooth structure. You can then continue to the low frequency components of the same upper left corner of the 2nd, 3rd inferior wavelet transform [6].

Saliva pH testing in predicting dental caries in children aged 7–10 years; Magdalena Fudali-Walczak , Grzegorz Raba, Beata Obłozą. 2015; thirteen (2):90-94. Dental caries stays the most everyday sickness of the masticatory system. is intently associated to modern-day dwelling conditions, its severity in particular issues populations of those countries where refined sugar is bump off in massive quantities. World agencies dealing with public fitness such as WHO or FDI are sounding alarm because, notwithstanding the formulated goals to be performed by using the year 2000, the epidemic of dental caries has now not been halted, or even, in mild of current research, has now not been reduced. The decay-inducing factor is dental bacterial plaque adhering to teeth, and the specific microorganisms it contains[4].

Correlation between the PH of saliva, plaque and buffering potential of saliva Dental caries is a disorder the place bacterial techniques cause harm to the hard teeth ,structure, characterized via acid demineralization (represented by pH) of the enamel enamel. Changes to the micro flowers within the oral cavity result in an overgrowth of more than a few bacteria like mutans streptococci and Lactobacilli that cause Dental Caries by means of producing acids. A find out about designed with 50 topics from whom prompted saliva samples and plaque have been amassed to measure the buffering capability with the use of chair-side kits (CRT Buffer, Ivoclar Vivadent AG, Schaan, Liechtenstein) and pH with a chair side test strip, Hydriion (9800) Spectral 0-14 Plastic pH Strip (Micro Essential Laboratories, USA).The records have been processed statistically with the Statistical Package for Social Sciences. [2].

Saliva as a Diagnostic Tool for Assessment of Dental Caries; V.K. Gopinath, A.R. Arzreanne School of Dental Sciences, The secretion charge and fine of saliva are necessary no longer only in caries development however also for remineralization. Saliva is indispensable to the integrity of the teeth as well as the soft tissues. It is now not unusual to take a look at patients



presenting with loss of enamel structures due to dental caries or dental erosion. The most feasible motives for exchange in the oral balance that is favoring demineralization may also be answered with the aid of measuring essential salivary parameters. Evaluating the causative elements in saliva of individual's at hazard to dental caries can pay the way to make suggestions that will cater specifically to individual's needs. Many benefits exist for each sufferers and dentist through introducing practice can advantage from improved diagnostics, early detection of problems, improved affected person communication and motivation and an increased dental attention for patients. Hence this learn about of salivary testing in dental caries assessment[3].

Human saliva is now not just the fluid in our mouth, however it mirrors our body's fitness and well being. Biomolecules that are circulating in the blood are additionally found in human saliva. It consists of approximately about 2,000 proteins, and most significantly, 26% of these proteins are additionally determined in blood, therefore emphasizes the saliva's significance as an introduced organic useful resource for ailment analysis and monitoring, as well as an remaining diagnostic medium to set up a person's response to treatment. The area of saliva diagnostics (SDs) started out in the early 60s when salivary calcium ranges were found to be extended in cystic fibrosis patients, and 50 years on now how the area has unmitigated to an unpredicted distance due to the improvement of increasingly touchy detection techniques. Dr. Shantala Arunkumar Dr. Arunkumar. J S Assistant Professor, Department of Oral Medicine and Radiology, SDM College of Dental Sciences and Hospital, Karnataka, India DWT Decomposition model[7].

salivary biomarkers help in early detection of diseases and increase the rate of success of the treatment. Sampling of saliva is safe, simple, cost effective and does not demand an expertise for collection. Therefore, saliva can be an important diagnostic armamentarium for mass screening for a specific disease or in remote areas. However, despite of various research, there are no definite guidelines regarding sensitivity and specificity of salivary diagnostic tools. Health care professionals must go hand in hand with government agencies to develop more research so that a general acceptance can be developed like that of traditional blood/urine analysis. Mishra A. Department of Dental Sciences, King George Medical University, Lucknow-226003, India[19].

Sharmila Baliga, Sangeeta Muglikar, Rahul Kale Saliva contains a variety of host defense factors. It influences calculus formation and periodontal disease. Different studies have been done to find exact correlation of salivary biomarkers with periodontal disease. With a multitude of biomarkers and complexities in their determination, the salivary pH may be tried to be used as a quick chairside test. The aim of this study was to analyze the pH of saliva and

determine its relevance to the severity of periodontal disease[20].

Saliva contains a lot of host defense factors. Wide arrays of studies have been done to locate genuine correlation of salivary markers with oral health status. With a multitude of biomarkers, the salivary pH may additionally be tried to be used as a rapid chair side take a look at for determining the oral health status. The aim of this study was once to analyze the pH of saliva and decide its relevance to the oral fitness status. The study population consisted of 60 patients. They had been divided into two groups of 30 patients each: Group A had clinically healthy gingiva, Group B who had grade -III calculus and generalized continual periodontitis. The randomized unstimulated saliva from each patient was once accumulated and pH used to be tested. Data was analyzed statistically the use of analysis of variance technique. The salivary pH was more acidic for sufferers with grade -III calculus and generalized chronic periodontitis as compared with the manage team ($P = 0.001$). These effects point out a vast change in the pH depending on the oral hygiene popularity and severity of the periodontal condition. The salivary pH indicates considerable modifications and for that reason relevance to the oral hygiene repute and severity of periodontal disease. Salivary pH may additionally therefore be used as a speedy chairside diagnostic biomarker of oral fitness status Dr. Saranya Manikandan, Ayyappan, T. Ilanchezhian, Vemulapalli Saimahitha[22].

Oral and dental diseases: Causes, prevention and treatment strategies naseem shaha. Dental caries is an infectious microbiological disease of the teeth that effects in localized dissolution and destruction of the calcified tissues. It is the 2d most common cause of teeth loss and is observed universally, irrespective of age, sex, caste, creed or geographic location. It is considered to be a disorder of civilized society, related to way of life factors, but heredity also performs a role. In the late stages, it causes severe pain, is luxurious to treat and leads to loss of precious man-hours. However, it is preventable to a certain extent. The incidence of dental caries in India is 50%–60%[12].

III. METHODOLOGY

3.1 Methods for collection of saliva per se:

Whole Saliva: The authors have for my part attempted various methods of collection total saliva, both stimulated and unstimulated conditions. Whole salivary collection is handy and non-invasive. It has been found to be finest and clinically more relevant in the assessment of overall salivary gland dysfunction. But, individual gland secretions are considered to be superior to whole saliva for many compositional analytes, because whole saliva carries non-salivary elements such as desquamated epithelial cells, bacteria, GCF and leukocytes.

The methods handy currently for collection of whole saliva include draining, spitting, suction and swab method. Shannon has in contrast flow prices in a group of subjects in special body positions. He suggested higher flow price values in the standing function and lower values in the lying role as compared with the go with the flow price in the sitting position.(9) Thus, it is perfect to collect saliva, while the difficulty is sitting upright with the head slightly tilted ahead and the eyes open.

Draining Method:

The problem is made to sit down quietly with the head bent down and the mouth open to permit the saliva to drip passively from the decrease lip into the graduated sterile tubes. Saliva amassed by way of draining is without any stimulation and is more reliable.



Fig.3. 1: Draining Method

Spitting Method: Saliva is allowed to accumulate in the floor of the mouth and the difficulty spits out it into the preweighed or graduated test tubes. The advantage of this approach is that it can be used when the glide price is very low and the place evaporation of saliva has to be minimised. The drawback is that it may have some stimulatory effect, and consequently can't be used for unstimulated saliva collection.



Fig.3. 2: Spitting Method

Suction Method: Saliva is allowed to accumulate in the floor of the mouth and aspirated always using micropipettes, syringes, saliva ejector or an aspirator.

Swabbing Method: It is carried out with the aid of introducing a synthetic gauze sponge, pre-weighed swab or cotton pad into the mouth, at the orifices of fundamental salivary glands. The topics are asked to chunk such that the sponge gets soaked within the saliva. Saliva soaked sponge is removed and placed in a sterile check tubes. Though this method is much less reliable, it helps in the evaluation of the level of oral dryness. It is in general used in the monitoring of drugs, hormones or steroids.


3.1.1 Dental diseases:

1 dental caries

Dental caries is one of the most common preventable diseases which is diagnosed as the most important cause of oral ache and enamel loss. Dental caries, a continual disease is unique amongst human and is one of the most common essential global oral health issues in the world today. It is the destruction of dental difficult a cellular tissue by means of acidic by-products from the bacterial fermentation of dietary carbohydrates mainly sucrose. It progresses slowly in most of the people which outcomes from an ecological imbalance in the equilibrium between teeth minerals and oral biofilms which is characterised by microbial activity, resulting in fluctuations in plaque pH due to bacterial acid production, buffering action from saliva and the surrounding enamel structure. The microbial community of caries is numerous and consists of many facultatively and obligately-anaerobic bacteria. *S. mutans* is the most principal associated with it. Dental caries can affect the human in quite a number methods i.e. presence of enamel pain, contamination or dysfunction of the stomatognathic device can limit the fundamental ingestion of energetic foods, affecting the boom in youngsters and adults as properly as their learning, verbal exchange skills and recreational things to do.

Types of caries:

- i) incipient caries/primary caries: Decay at a vicinity that has not skilled preceding decay.
- ii) recurrent caries/secondary caries: Appears at a region with a preceding history of caries and is often discovered on the margins of fillings and other dental restorations.
- iii) arrested caries: A lesion on a teeth that used to be in the past demineralized but was remineralized earlier than causing a cavitation.[6]



3.3 Caries aetiology

The recent year children's oral health outcomes using a border frame work, which incorporates psychological and environmental predictors as well as the biological and dietary effect. This framework generally classifies conditions associated with disease into five broad domains: genetics and biology, social envormental, physical, environment, health influencing behaviour and medical care.

Child level: visible plaque early colonization by carries related bacteria ,the presence of mutons streptococci ,frequent intake of sweetened drinks, in frequent tooth brushing, in less and use of antibiotics have all been associated with carries development in preschool children's [9]

Family level:

It is associated with carries risk in children included are demographic factor of the family, parental oral heath behavior and attitude, dental anxiety and dental attendance. Maternal health and life style in pregnancy and early childhood. Community it is associated with carries risk in children included are demographic factor of the family, parental oral heath behavior and attitude, dental anxiety and dental attendance .maternal health and life style in pregnancy and early childhood. [9]

Community level:

Children's oral health is lightly to be better in a community that values good oral health cultural aspect as neighborhood may have implications for carries development.

2 periodontitis:

The essential constituent of this hypotonic biofluid is water comparising 99.5% of the whole volume, with the relaxation of 0.5% being amino acids, histatins, cystatins, defensins, statherins, lysozyme, proline-rich proteins, carbonic anhydrases, peroxidases, lactoferrin, mucins, secretory immunoglobulins, lipids, together with various ions, such as potassium, calcium, chloride, sodium and phosphates. The

foundation of chemical and biochemical aspects of salivary fluid is numerous and complex, and these are no longer covered in the current review. Periodontal troubles are among the most essential and frequent adult's gum disorders. The severity of periodontitis ought to be diagnosed on the groundwork of its typical clinical parameters. These include periodontal probing pocket depth, clinical attachment and amount of gum bleeding. Analysis of saliva as an important laboratory test for the assessment of many salivary conditions, including periodontitis, has attracted periodontitis' attention at some point of the closing few a long time. Saliva incorporates each regionally and systemically derived biochemical's with relatively necessary diagnostic value, which could be used for detecting periodontal disorders. They are many times used to perceive the type and severity of periodontitis, as well as monitoring the remedy success. A range of advanced methods are regarded nowadays which use saliva for diagnosis of some oral and systemic diseases. The early-stages of periodontitis is recognized as gingivitis and, in most cases, it has no signs and symptoms and is now not painful. Therefore, it is very important with clinical benefits to diagnose the ailment at early tiers the use of a sensitive and reliable method. Detection of salivary biomarkers is a non-invasive laboratory examination for early diagnosis of periodontitis[11].

Respiratory disease is one of the most common causes of mortality in adults which may also contaminate the oral cavity, inflicting periodontal problems. Human oral cavity is closed to the trachea and any respiratory pathogen passes via it. Therefore, the formation of dental plaque need to be avoided, in order to reduce the risk of contamination in the respiratory tract. It has been proven that persistent obstructive pulmonary disease is not solely related to smoking, however it additionally has a relationship with periodontitis.

Cardiovascular disease might also somehow be associated to a variety of sorts of periodontal problems. This truth has been confirmed for a crew of young men. Alveolar bone and attachment loss (indicators of periodontitis) have also been reported with the aid of radiographic examination . It has been proven that young men suffering from periodontal troubles are at greater threat of stroke than sufferers in the same age crew with moderate or no periodontitis . However, bad oral health influences systemic health, given that many ordinary chance factors for cardiovascular disease and stroke (e.g., smoking, being male, sedentary lifestyle, obesity, hypertension and dyslipidemia) are hard to change, while improvement in oral care might effortlessly be modified[11].

3.2 Methods of Salivary Test

The point so interest of two spit two examining making use of pH sensor for discovering verbal illnesses. The most important advantage of salivary take a look at over chemical blood test is



its security. The make use of of needles in blood collection making use of cannot be completely free of chance from sickness inflicting infections. Since salivary test can also be a noninvasive procedure, at that factor sufferers who are us of two now confined from utilizing needles, such as teens ,the matured, and hemophiliac patients, will be managed a quantity of preferences in self-monitoring. Next, not at all like blood, saliva can be self-collected, and no longer at all like pee, spit can be amassed any time. These are the fundamental conditions that POCT need to meet in arrange to structure on-site trying out feasible at domestic and within the working environment. More over, it is mainly treasured for mass screening tests. On the groundwork of the performed check-ups of the oral cavity, a DMF index was once hooked up for every patient. The saliva pH check was carried out in the following way: a vial of round 10millilitre of inspired saliva, The affected person is knowledgeable in strengthen no longer to take any food or drink two hours prior to the test. The end result of the check used to be examine by means of ability of a pH meter two after 30 seconds from dipping the pH sensor in saliva. The consequences received were statistically analyzed by means of potential of the arduino. Tabular descriptions have been used in the research – facts is presented in the form of tables together with the relationships of the most considerable research hypotheses; the cross tabs include basic impartial variables. The goal of prophylactic programmers conducted by means of dentists is patient training in the region of prevention of the ailments of the oral cavity, which to a wonderful diploma relies upon on oral cavity hygiene, diet, dental hygiene visits and treatment in dental practice. Information regarding cleanliness of the verbal despair are an necessary two cleanliness of the two verbal despair are an necessary indicator for the persistent and a huge thought to improve prohealth exercises. They permit the expert to draw sufferers consideration to performing getting ready on hygiene of the verbal depression, as well as recognizing a extra considerable no. of patients, human beings at greater hazard for rot. The investigation has proved that spit takes portion within the take care of of mineralization and demineralization of the difficult enamel tissues. At pH levels between 6.8 and 7.2 spit should be a supersaturated arrangement of calcium phosphate;hence after a slite demineralization misplaced minerals factors can return to the challenging tissues of the tooth from spit.

Saliva has a cleaning impact on the teeth. Normally, 700–800 ml of saliva is secreted per day. Caries recreation increases as the viscosity of the saliva increases. Eating fibrous food and chewing vigorously will increase salivation, which helps in digestion as well as improves cleaning of the teeth. The quantity as properly as composition, pH, viscosity and buffering capacity of the saliva plays a position in dental caries.

- Quantity: Reduced salivary secretion as determined in xerostomia and salivary gland a plasia offers upward jostle to expanded caries activity.

- Composition: Inorganic—fluoride, chloride, sodium, magnesium, potassium, iron, calcium and phosphorus are inversely related to caries. Organic—ammonia retards plaque formation and neutralizes the acid.
- pH: A impartial or alkaline pH can neutralize acids formed by the motion of microorganisms on carbohydrate food substances.
- Antibacterial factors: Saliva incorporates enzymes such as lactoperoxidase, lysozyme, lactoferrin and immunoglobulin (Ig)A, which can inhibit plaque bacteria.




Figure 3.4 flowchart of pathogenesis of dental of dental caries

The sucrose connection related to caries Sucrose is a fundamental environmental contributor to dental caries because of the many oral streptococci possess extracellular enzyme succesful of cleaving the α -1 and α -2 glycosidic bond of sucrose and harnessing the electricity to yield glucose polymer (glucans and mutans) and fructose. This crew of enzymes referred to as the glucosyltransferases (GTFs), money owed for this unique relationship between sucrose and caries. Formation of glucan-mutan polymer allows the cariogenic bacteria to accumulate into biofilm to form a essential mass. Without formation of crucial mass cariogenic micro organism would colonize the oral cavity but no longer be massed so as to cause the destruction of the enamel surface. Accordingly the formation of essential mass that is uniquely related with glucosyltransferases and sucrose is a organic motive for man’s recent suffering with caries[6].



IV. CONCLUSION

This find out about confirmed that salivary parameters such as salivary go with the flow rate, salivary viscosity, salivary. pH and salivary buffering potential have been decrease in subjects with high dental caries. Hence, we recommend salivary trying out to be a part of routine prognosis when treating patient with high dental caries risk. Dental caries remains a extensively common bacterial infection regardless of terrific advances in prevention and treatment, and continues to incorporate a huge element of whole U.S. expenditures on fitness care. Why caries continues to be a essential public fitness hassle remains an unanswered question, however perception may. Be won via evaluation of the danger elements associated with the disease. This file has evaluated the proof implicating salivary parameters as possibly chance factors. Saliva is a complicated body fluid that presents a generic shielding characteristic for exposed oral tough tissues. Several medical conditions, such as Sjögren's syndrome, therapeutic radiation to the head and neck, and pharmacological retailers with xerostomic aspect effects, decrease salivary float charge to pathological degrees and dramatically raise the patient's risk of caries. Buffering capacity of saliva reduces the pH of saliva to some extent, but has not a great deal function in the discount of pH of plaque. There exists a need to discover tools which can help to define the diploma of decay hazard for a given patient, as properly as elements motivating the patient to undertake actions enhancing the situation of hygiene in the oral cavity. Data concerning hygiene of the oral cavity are an important indicator for the patient and a huge motivation to improve prohealth activities. They enable the health practitioner to draw patient's interest to performing training on hygiene Of the oral cavity, as properly as figuring out a wider team of patients, people at higher chance for decay. The imaging of the situation of oral cavity hygiene is an important issue of health schooling in dentistry. It's important purpose is making two the lookup population aware of the significance of prophylaxis of the ailments of the oral cavity. The purpose of prophylactic programs carried out by dentists is patient training in the place of prevention of the ailments of the oral cavity, which to a exquisite diploma relies upon on oral cavity hygiene, diet, dental hygiene visits and remedy in dental practice. Taken together saliva has a tremendous possible of becoming the next prognosis fluid of choice due to functional correlations that can be made between salivary markers and distinctive diseases. However more studies are wished in order to identify specific biomarkers or panels of biomarkers that can be used for diagnosis and monitoring in medical settings.

B. FUTURE SCOPE:

Salivary diagnostics may want to dramatically alternate scientific exercise by means of way of introducing point of care checking out and real-time by using disorder surveillance. Dentistry can help this imaginative and prescient quit up

reality, in area due to the fact many sufferers see their dentist greater often in part be than their health practitioner because of the prevention focal factor and the need for ordinary visits to keep oral health. While fantastically inexpensive, non-invasive simple and accurate/diagnostics take a seem at techniques provide obvious, benefits, there are also challenges and distinct considerations that are additional associated with their activities use two two in dental practice. Importantly the dental career need to address know-how gaps that may additionally exist. Involving now not solely the pre-requisites of which tests are designed, then again additionally a differ of economic (reimbursement) and a vary of inter-professional practice implications. Although the dental workplace can be a important putting for the implementation of oral fluid tests, some can also additionally feel that embracing a broader base of clinical diagnostics, such as breast most cancers screening or drug monitoring, as a alternative than a focal point on checks for oral prerequisites (e.g., dental caries and periodontal disease), can also no longer be notable for their practice.

V. REFERENCE

- [1] K..Yamuna Priya, K.muthu prathibha "Method of collection saliva a review" International journal of oral health dentistry DOI: 10.18231/2395-499X.2017.0032 july-september 2017.
- [2] B.Vijaya lakshmi, surendra reddy,munnangi,manoj kumar,"corrrelation between the ph of saliva,plaque nad buffering capacity of saliva" international journal of applied dental science pp;48-50september2017.
- [3] V.K.gopinath,A.R.arzreanne"saliva as a diagnostics tool for assessment of dental caries",archieve of orofacial sciences pp; 1: 57-59,2006.
- [4] W.Magdalena fudali ,grzegorz raba,beata obloza, "saliva pH testing in predicting dental caries in children aged 7-10 years," Vision, Image and Signal Processing, IEE Proceedings -, vol. 152, pp. 561-574, 2005.
- [5] Ying sun,wei du,chunguang zhou, " A computational method for prediction of saliva secretory proteins and its application to indentification of head and neck cancer biomarkers for saliva diagnosis ", In IEEE transaction on nanobioscience,vol,no.14,2march2015.
- [6] Yadav khushbu and Satyam Prakash. Asian Journal of Biomedical and Pharmaceutical Sciences, 6(53), 2016, 01-07.doi: 10.15272/ajbps.v6i53.773, 2016.
- [7] Shantala anilkumar,krishna n burde" development in diagnostics application of saliva in oral and systematic disease-a comprehensive review", Journal of Scientific and Innovative Research 2014; 3(3): 372-387



- [8] Cataldo W. Leone, D.M.D., D.M.Sc.; Frank G. Oppenheim, D.M.D., Ph.D. "Physical and chemical aspects of saliva as indicator of risk for dental caries in humans", *Journal of dental education* vol.65,no.10.
- [9] William v.giannobile ,thomas beikler,janet s.kinney,"Saliva as a diagnostic tool for periodontal disease:current state and future directions.", *periodontology* 2000,vol.50,2009,pp-52-64.
- [10] Jooana cunha cruz dds,joanna scott, "salivary characteristics and dental carries" [http:// jada. ada. Org .](http://jada.ada.org) *JADA* 2013;144(5):e31-e40.may2013.
- [11] Mojgan paknjad, afsaneh rezaei"salivary biochemical biomarkers of periodontitis", *dental research center,department of periodontology april ROM. J. BIOCHEM.*, 50, 2, pp-129-146 (2013)12,2013.
- [12] Shah naseem , "oral and dental diseases:causes, prevention and treatment strategies", nov2003.pp-275-298,2003.
- [13] Radha s.p.malon,sahba sadir,malarvilli balakrishnan, "saliva based biosensors:noninvasive monitoring tool for clinical diagnostics", *biomed research*, Vol. 2014, No. 12, pp. 20, september, 2014.
- [14] Bradley Henson, DDS, PhD Ronald Zentz, RPh, DDS David T. Wong, DMD, DMSc, " A Primer on Salivary Diagnostics ", in *Proceedings of the International Conference on Multimedia and Expo*, pp. 429-432, 2001.
- [15] Chitharanjan M Shetty, Mithra Nidarsh Hegde " Correlation between dental caries with salivary flow, pH, and buffering capacity in adult south Indian population: An in-vivo study", *IEEE Transactions on Circuits and Systems- II*, Vol. 45, No. 8, pp. 1097-1101, August 1998.
- [16] Maria Greabu* and Bogdan Calenicl, " Saliva: A Diagnosis Fluid for Oral and General Diseases ", Greabu and Calenic, *Organic Chem Curr Res* 2015, 4:1,DOI: [10.4172/2161-0401.1000131](https://doi.org/10.4172/2161-0401.1000131).
- [17] Kashi Raj Bhattarai(2018), Hyung-Ryong Kim, Han-Jung Chae" Compliance with Saliva Collection Protocol in Healthy Volunteers: Strategies for Managing Risk and Errors" *Int. J. Med. Sci.*, Vol. 15 2018; 15(8): 823-831. doi: 10.7150/ijms.25146.
- [18] Vivek Vardhan Gupta, Neha Chitkara, Harsh Vardhan Gupta, Arshdeep Singh, Ramandeep Singh Gambhir, Harkiranjot Kaur" Comparison of Salivary Calcium Level and pH in patients with Aggressive Periodontitis and Healthy Individuals: A Clinico -Biochemical Study" *OHDM - Vol. 15 - No. 2 -pp-122-126*, April, 2016.
- [19] Mishra A" Saliva Biomarkers: The Unsung Hero of Diagnostics" *J Mol Genet Med*, an open access journal Volume 11 • Issue 3 • DOI: [10.4172/1747-0862.1000288](https://doi.org/10.4172/1747-0862.1000288).
- [20] Sharmila Baliga, Sangeeta Muglikar, Rahul Kale" Salivary pH: A diagnostic biomarker" Department of Periodontology and Implantology,M.A. Rangoonwala College of Dental Sciences and Research Centre, Pune,Maharashtra,IndiaDOI:10.4103/0972-124X.118317. April 3, 2018.
- [21] Chen-Zi Zhang, Xing-Qun Cheng"Saliva in the diagnosis of diseases" *International Journal of Oral Science* (2016) 8,pp- 133-137,
- [22] Manikandan, Dr. Saranya , Ayyappan, T.Ilanchezian., Vemulapalli Saimahitha" Salivary pH as a Diagnostic Marker of Oral Health Status" *Int. J. Pharm. Sci. Rev. Res.*, 51(1), July - August 2018; Article No. 07, Pages: 41-44.